

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
20 January 2005 (20.01.2005)

PCT

(10) International Publication Number
WO 2005/005374 A1

(51) International Patent Classification⁷: **C07C 237/22**,
237/12, 237/14, C07D 333/02, 307/02, 211/04, 249/08,
C07C 311/16, 311/08, C07D 275/02, 285/06, 213/04,
A61K 31/166, A61P 25/28, C07K 5/02, C07D 241/08

(72) Inventor: YANG, Wenjin; 383 Bodega Street, Foster City,
CA 94404 (US).

(21) International Application Number:

PCT/US2003/039276

(22) International Filing Date:

10 December 2003 (10.12.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

10/462,127

16 June 2003 (16.06.2003)

US

PCT/US03/18858

16 June 2003 (16.06.2003)

US

(71) Applicant: **SUNESIS PHARMACEUTICALS, INC.**
[US/US]; 341 Oyster Point Boulevard, South San Fran-
cisco, CA 94080 (US).




(74) Agent: LAGNEAU, Nadège, M.; Choate, Hall & Stewart,
Exchange Place, 53 State Street, Boston, MA 02109 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ,
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,

[Continued on next page]

(54) Title: ASPARTYL PROTEASE INHIBITORS



(57) Abstract: The present invention provides compounds having the formula: (I) wherein R^1 , R' , R^2 , R^3 , R^4 , X^1 , X^2 and X^3 are as defined herein, and pharmaceutical compositions thereof. The present invention also provides methods of inhibiting proteases, more specifically aspartyl proteases. In certain embodiments, compounds inhibit BACE (β -site APP-cleaving enzyme), and thus are useful in the treatment or prevention of a disease characterized by β -amyloid deposits in the brain (including, but not limited to, Alzheimer's Disease). The present invention also provides methods for preparing compounds of the invention.



SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, — *with amended claims*
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

ASPARTYL PROTEASE INHIBITORS

PRIORITY CLAIM

[0001] The present application claims priority to U.S. Patent Application No.: 10/462,127; filed June 16, 2003, and Patent Cooperation Treaty Application No.: PCT/US03/18858, filed June 16, 2003; each of which claims priority to U.S. Provisional Patent Application Nos.: 60/430,693, filed December 3, 2002, and 60/389,194, filed June 17, 2002. The entire contents of each of the above-referenced applications are incorporated herein by reference.

BACKGROUND OF THE INVENTION

[0002] Alzheimer's Disease is a progressive dementia in which massive deposits of aggregated protein breakdown products (β -amyloid plaques and neurofibrillary tangles) accumulate in the brain, resulting in the loss of memory, cognition, reasoning, judgement, orientation, and eventually death. Current therapies for the treatment of Alzheimer's Disease include, but are not limited to, donepezil and tacrine. These therapies are useful for improving the memory of patients during the early stages of Alzheimer's Disease, however they do not modify the progression of aggregated protein breakdown products underlying the pathology of Alzheimer's Disease. It would be desirable to develop therapies that would either stop or slow down this process of aggregation.


[0003] As described above, a defining feature of Alzheimer's Disease which is often used during clinical diagnosis is the presence of β -amyloid plaques and neurofibrillary tangles. β -amyloid plaques are predominantly composed of amyloid β peptide ($A\beta$ (or $\beta A4$), which is derived by proteolysis of the amyloid precursor protein (APP). Proteolysis of the amyloid precursor protein is effected by several enzymes called secretases. More specifically, cleavage of APP at the N-terminus of the $A\beta$ peptide by β -secretase and at the C-terminus by one or more γ -secretases constitutes the β -amyloidogenic pathway, *i.e.*, the pathway by which $A\beta$ is formed. It is believed that $A\beta$ peptide accumulates as a result of this APP processing by β -secretase and thus inhibition of this enzyme's activity is desirable for the treatment of Alzheimer's Disease. For example, *in vivo* processing of APP at the β -secretase

cleavage site is thought to be a rate limiting step in A β production, and is thus believed to be a therapeutic target for Alzheimer's Disease (Sabbagh *et al. Alz. Dis. Rev.* 1997, 3, 1-19). Recently, an aspartyl protease (known as BACE, Asp2, Memapsin) has been identified as the enzyme responsible for processing of APP at the β -secretase cleavage site (see, for example, Vassar, *et al. Science*, 1999, 286, 735-741; Yan *et al. Nature*, 1999, 402, 533-537; Sinha *et al. Nature*, 1999, 402, 537-540; and Hussain *et al. Mol. Cell. Neurosci.* 1999, 14, 419-427).

[0004] Because it is believed that BACE plays an important role in the development and pathogenesis of Alzheimer's Disease, there has been increasing interest in the development of inhibitors of BACE as treatments (and possibly as preventative agents) for Alzheimer's Disease and other disorders caused by the accumulation of β -amyloid plaques. There remains a need, however, for the development of novel therapeutics capable of inhibiting the activity of this aspartyl protease. In particular, it would be desirable to develop therapeutics capable of selectively inhibiting BACE.

SUMMARY OF THE INVENTION

[0005] As discussed above, there remains a need for the development of novel therapeutic agents and agents useful for treating disorders mediated by aspartyl proteases. The present invention provides novel compounds having the structure:



(I)

and pharmaceutical compositions thereof, as described generally and in subclasses herein, which compounds are useful as inhibitors of aspartyl proteases, and thus are useful, for example, for the treatment of Alzheimer's Disease.

[0006] In certain other embodiments, the invention provides pharmaceutical compositions comprising an inventive compound, wherein the compound is present in an amount effective to inhibit β -secretase activity. In certain other embodiments, the invention provides pharmaceutical compositions comprising an inventive compound and optionally further comprising an additional therapeutic agent. In yet

other embodiments, the additional therapeutic agent is an agent for the treatment of Alzheimer's Disease.

[0007] In yet another aspect, the present invention provides methods for inhibiting β -secretase activity in a patient or a biological sample, comprising administering to said patient, or contacting said biological sample with an effective inhibitory amount of a compound of the invention. In still another aspect, the present invention provides methods for treating any disorder involving β -secretase activity, comprising administering to a subject in need thereof a therapeutically effective amount of a compound of the invention. In certain other embodiments, the invention provides a method for treating or preventing a disease characterized by β -amyloid deposits in the brain comprising administering to a patient a therapeutically effective amount of a compound of the invention.

BRIEF DESCRIPTION OF THE DRAWING

[0008] Figure 1A depicts a plasma concentration curve for an exemplary inventive compound.

[0009] Figure 1B depicts a brain concentration curve for an exemplary inventive compound.

DEFINITIONS

[0010] Certain compounds of the present invention, and definitions of specific functional groups are described in more detail below. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed., inside cover, and specific functional groups are generally defined as described therein. Additionally, general principles of organic chemistry, as well as specific functional moieties and reactivity, are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999, the entire contents of which are incorporated herein by reference. Furthermore, it will be appreciated by one of ordinary skill in the art that the synthetic methods, as described herein, utilize a variety of protecting groups. By the term "protecting group", as used herein, it is meant that a particular functional moiety, *e.g.*, O, S, or N, is temporarily blocked so

that a reaction can be carried out selectively at another reactive site in a multifunctional compound. In preferred embodiments, a protecting group reacts selectively in good yield to give a protected substrate that is stable to the projected reactions; the protecting group must be selectively removed in good yield by readily available, preferably nontoxic reagents that do not attack the other functional groups; the protecting group may form an easily separable derivative (more preferably without the generation of new stereogenic centers); and the protecting group has a minimum of additional functionality to avoid further sites of reaction. As detailed herein, oxygen, sulfur, nitrogen and carbon protecting groups may be utilized. For example, in certain embodiments, as detailed herein, certain exemplary oxygen protecting groups are utilized. These oxygen protecting groups include, but are not limited to methyl ethers, substituted methyl ethers (*e.g.*, MOM (methoxymethyl ether), MTM (methylthiomethyl ether), BOM (benzyloxymethyl ether), PMBM (p-methoxybenzyloxymethyl ether), to name a few), substituted ethyl ethers, substituted benzyl ethers, silyl ethers (*e.g.*, TMS (trimethylsilyl ether), TES (triethylsilylether), TIPS (triisopropylsilyl ether), TBDMS (t-butyldimethylsilyl ether), tribenzyl silyl ether, TBDPS (t-butyldiphenyl silyl ether), to name a few), esters (*e.g.*, formate, acetate, benzoate (Bz), trifluoroacetate, dichloroacetate, to name a few), carbonates, cyclic acetals and ketals. In certain other exemplary embodiments, nitrogen protecting groups are utilized. These nitrogen protecting groups include, but are not limited to, carbamates (including methyl, ethyl and substituted ethyl carbamates (*e.g.*, Troc), to name a few) amides, cyclic imide derivatives, N-Alkyl and N-Aryl amines, imine derivatives, and enamine derivatives, to name a few. Certain other exemplary protecting groups are detailed herein, however, it will be appreciated that the present invention is not intended to be limited to these protecting groups; rather, a variety of additional equivalent protecting groups can be readily identified using the above criteria and utilized in the present invention. Additionally, a variety of protecting groups are described in "Protective Groups in Organic Synthesis" Third Ed. Greene, T.W. and Wuts, P.G., Eds., John Wiley & Sons, New York: 1999, the entire contents of which are hereby incorporated by reference

[0011] It will be appreciated that the compounds, as described herein, may be substituted with any number of substituents or functional moieties. In general, the term “substituted” whether preceded by the term “optionally” or not, and substituents contained in formulas of this invention, refer to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. When more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. As used herein, the term “substituted” is contemplated to include all permissible substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and non-aromatic substituents of organic compounds. For purposes of this invention, heteroatoms such as nitrogen may have hydrogen substituents and/or any permissible substituents of organic compounds described herein which satisfy the valencies of the heteroatoms. Furthermore, this invention is not intended to be limited in any manner by the permissible substituents of organic compounds. Combinations of substituents and variables envisioned by this invention are preferably those that result in the formation of stable compounds useful in the treatment and prevention, for example of disorders, as described generally above. The term “stable”, as used herein, preferably refers to compounds which possess stability sufficient to allow manufacture and which maintain the integrity of the compound for a sufficient period of time to be detected and preferably for a sufficient period of time to be useful for the purposes detailed herein.

[0012] The term “aliphatic”, as used herein, includes both saturated and unsaturated, straight chain (*i.e.*, unbranched), branched, cyclic, or polycyclic aliphatic hydrocarbons, which are optionally substituted with one or more functional groups. As will be appreciated by one of ordinary skill in the art, “aliphatic” is intended herein to include, but is not limited to, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, and cycloalkynyl moieties. Thus, as used herein, the term “alkyl” includes straight, branched and cyclic alkyl groups. An analogous convention applies to other generic terms such as “alkenyl”, “alkynyl” and the like. Furthermore, as used herein, the terms “alkyl”, “alkenyl”, “alkynyl” and the like

encompass both substituted and unsubstituted groups. In certain embodiments, as used herein, "lower alkyl" is used to indicate those alkyl groups (cyclic, acyclic, substituted, unsubstituted, branched or unbranched) having 1-6 carbon atoms.

[0013] In certain embodiments, the alkyl, alkenyl and alkynyl groups employed in the invention contain 1-20 aliphatic carbon atoms. In certain other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-10 aliphatic carbon atoms. In yet other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-8 aliphatic carbon atoms. In still other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-6 aliphatic carbon atoms. In yet other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-4 carbon atoms. Illustrative aliphatic groups thus include, but are not limited to, for example, methyl, ethyl, n-propyl, isopropyl, cyclopropyl, -CH₂-cyclopropyl, allyl, n-butyl, sec-butyl, isobutyl, tert-butyl, cyclobutyl, -CH₂-cyclobutyl, n-pentyl, sec-pentyl, isopentyl, tert-pentyl, cyclopentyl, -CH₂-cyclopentyl-n, hexyl, sec-hexyl, cyclohexyl, -CH₂-cyclohexyl moieties and the like, which again, may bear one or more substituents. Alkenyl groups include, but are not limited to, for example, ethenyl, propenyl, butenyl, 1-methyl-2-buten-1-yl, and the like. Representative alkynyl groups include, but are not limited to, ethynyl, 2-propynyl (propargyl), 1-propynyl and the like.

[0014] The term "alkoxy" (or "alkyloxy"), or "thioalkyl" as used herein refers to an alkyl group, as previously defined, attached to the parent molecular moiety through an oxygen atom or through a sulfur atom. In certain embodiments, the alkyl group contains 1-20 aliphatic carbon atoms. In certain other embodiments, the alkyl group contains 1-10 aliphatic carbon atoms. In yet other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-8 aliphatic carbon atoms. In still other embodiments, the alkyl group contains 1-6 aliphatic carbon atoms. In yet other embodiments, the alkyl group contains 1-4 aliphatic carbon atoms. Examples of alkoxy, include but are not limited to, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, tert-butoxy, neopentoxy and n-hexoxy. Examples of thioalkyl include, but are not limited to, methylthio, ethylthio, propylthio, isopropylthio, n-butylthio, and the like.

[0015] The term “alkylamino” refers to a group having the structure -NHR' wherein R' is alkyl, as defined herein. The term “dialkylamino” refers to a group having the structure -N(R')₂, wherein R' is alkyl, as defined herein. The term “aminoalkyl” refers to a group having the structure NH₂R'-, wherein R' is alkyl, as defined herein. In certain embodiments, the alkyl group contains 1-20 aliphatic carbon atoms. In certain other embodiments, the alkyl group contains 1-10 aliphatic carbon atoms. In yet other embodiments, the alkyl, alkenyl, and alkynyl groups employed in the invention contain 1-8 aliphatic carbon atoms. In still other embodiments, the alkyl group contains 1-6 aliphatic carbon atoms. In yet other embodiments, the alkyl group contains 1-4 aliphatic carbon atoms. Examples of alkylamino include, but are not limited to, methylamino, ethylamino, isopropylamino and the like.

[0016] Some examples of substituents of the above-described aliphatic (and other) moieties of compounds of the invention include, but are not limited to aliphatic; heteroaliphatic; aryl; heteroaryl; alkylaryl; alkylheteroaryl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; F; Cl; Br; I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)R_x; -S(O)₂R_x; -NR_x(CO)R_x; -N(R_x)CO₂R_x; -N(R_x)S(O)₂R_x; -N(R_x)C(=O)N(R_x)₂; -S(O)₂N(R_x)₂; wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, heteroaliphatic, aryl, heteroaryl, alkylaryl, or alkylheteroaryl, wherein any of the aliphatic, heteroaliphatic, alkylaryl, or alkylheteroaryl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic, and wherein any of the aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl substituents described above and herein may be substituted or unsubstituted. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein.

[0017] In general, the terms “aromatic moiety” and “heteroaromatic moiety”, as used herein, refer to stable mono- or polycyclic, heterocyclic, polycyclic, and polyheterocyclic unsaturated moieties having preferably 3-14 carbon atoms, each of which may be substituted or unsubstituted. It will also be appreciated that aromatic

and heteroaromatic moieties, as defined herein may be attached via an alkyl or heteroalkyl moiety and thus also include $-(\text{alkyl})\text{aromatic}$, $-(\text{heteroalkyl})\text{aromatic}$, $-(\text{heteroalkyl})\text{heteroaromatic}$, and $-(\text{heteroalkyl})\text{heteroaromatic}$ moieties. Thus, as used herein, the phrases "aromatic or heteroaromatic moieties" and "aromatic, heteroaromatic, $-(\text{alkyl})\text{aromatic}$, $-(\text{heteroalkyl})\text{aromatic}$, $-(\text{heteroalkyl})\text{heteroaromatic}$, and $-(\text{heteroalkyl})\text{heteroaromatic}$ " are interchangeable. Substituents include, but are not limited to, any of the previously mentioned substituents, *i.e.*, the substituents recited for aliphatic moieties, or for other moieties as disclosed herein, resulting in the formation of a stable compound.

[0018] The term "aryl", as used herein, does not differ significantly from the common meaning of the term in the art, and refers to a mono- or bicyclic carbocyclic ring system having one or two aromatic rings including, but not limited to, phenyl, naphthyl, tetrahydronaphthyl, indanyl, indenyl and the like.

[0019] The term "heteroaryl", as used herein, does not differ significantly from the common meaning of the term in the art, and refers to a cyclic aromatic radical having from five to ten ring atoms of which one ring atom is selected from S, O and N; zero, one or two ring atoms are additional heteroatoms independently selected from S, O and N; and the remaining ring atoms are carbon, the radical being joined to the rest of the molecule via any of the ring atoms, such as, for example, pyridyl, pyrazinyl, pyrimidinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, quinolinyl, isoquinolinyl, and the like.

[0020] It will be appreciated that aryl and heteroaryl groups (including bicyclic aryl groups) can be unsubstituted or substituted, wherein substitution includes replacement of one or more of the hydrogen atoms thereon independently with any one or more of the following moieties including, but not limited to: aliphatic; heteroaliphatic; aryl; heteroaryl; alkylaryl; alkylheteroaryl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; F; Cl; Br; I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)R_x; -S(O)₂R_x; -NR_x(CO)R_x; -N(R_x)CO₂R_x; -N(R_x)S(O)₂R_x; -N(R_x)C(=O)N(R_x)₂; -S(O)₂N(R_x)₂; wherein each occurrence of R_x

independently includes, but is not limited to, aliphatic, heteroaliphatic, aryl, heteroaryl, alkylaryl, or alkylheteroaryl, wherein any of the aliphatic, heteroaliphatic, alkylaryl, or alkylheteroaryl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic, and wherein any of the aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl substituents described above and herein may be substituted or unsubstituted. Additionally, it will be appreciated, that any two adjacent groups taken together may represent a 4, 5, 6, or 7-membered cyclic, substituted or unsubstituted aliphatic or heteroaliphatic moiety. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein.

[0021] The term "cycloalkyl", as used herein, refers specifically to cyclic moieties having three to seven, preferably three to ten carbon atoms. Suitable cycloalkyls include, but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like, which, as in the case of other aliphatic, heteroaliphatic or heterocyclic moieties, may optionally be substituted with substituents including, but not limited to aliphatic; heteroaliphatic; aryl; heteroaryl; alkylaryl; alkylheteroaryl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; F; Cl; Br; I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)R_x; -S(O)₂R_x; -NR_x(CO)R_x; -N(R_x)CO₂R_x; -N(R_x)S(O)₂R_x; -N(R_x)C(=O)N(R_x)₂; -S(O)₂N(R_x)₂; wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, heteroaliphatic, aryl, heteroaryl, alkylaryl, or alkylheteroaryl, wherein any of the aliphatic, heteroaliphatic, alkylaryl, or alkylheteroaryl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic, and wherein any of the aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl substituents described above and herein may be substituted or unsubstituted. Additionally, it will be appreciated that any of the cycloaliphatic or cycloheteroaliphatic moieties described above and herein may comprise an aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl moiety fused thereto. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein.

[0022] The term “heteroaliphatic”, as used herein, refers to aliphatic moieties in which one or more carbon atoms in the main chain have been substituted with a heteroatom. Thus, a heteroaliphatic group refers to an aliphatic chain which contains one or more oxygen sulfur, nitrogen, phosphorus or silicon atoms, *e.g.*, in place of carbon atoms. Heteroaliphatic moieties may be branched, unbranched, cyclic or acyclic and include saturated and unsaturated heterocycles such as morpholino, pyrrolidinyl, etc. In certain embodiments, heteroaliphatic moieties are substituted by independent replacement of one or more of the hydrogen atoms thereon with one or more moieties including, but not limited to aliphatic; heteroaliphatic; aryl; heteroaryl; alkylaryl; alkylheteroaryl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; F; Cl; Br; I; -OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)R_x; -S(O)₂R_x; -NR_x(CO)R_x; -N(R_x)CO₂R_x; -N(R_x)S(O)₂R_x; -N(R_x)C(=O)N(R_x)₂; -S(O)₂N(R_x)₂; wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, heteroaliphatic, aryl, heteroaryl, alkylaryl, or alkylheteroaryl, wherein any of the aliphatic, heteroaliphatic, alkylaryl, or alkylheteroaryl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic, and wherein any of the aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl substituents described above and herein may be substituted or unsubstituted.

[0023] In general, the term “cycloaliphatic”, as used herein, refer to a cyclic aliphatic moiety, wherein the term *aliphatic* is as defined above. A cycloaliphatic moiety may be substituted or unsubstituted and saturated or unsaturated. Substituents include, but are not limited to, any of the previously mentioned substituents, *i.e.*, the substituents recited for aliphatic moieties, or for other moieties as disclosed herein, resulting in the formation of a stable compound. In certain embodiments, cycloaliphatic compounds include but are not limited to monocyclic, or polycyclic aliphatic hydrocarbons and bridged cycloalkyl compounds, which are optionally substituted with one or more functional groups. As will be appreciated by one of ordinary skill in the art, “cycloaliphatic” is intended herein to include, but is not limited to, cycloalkyl, cycloalkenyl, and cycloalkynyl moieties, which are

optionally substituted with one or more functional groups. Illustrative cycloaliphatic groups thus include, but are not limited to, for example, cyclopropyl, -CH₂-cyclopropyl, cyclobutyl, -CH₂-cyclobutyl, cyclopentyl, -CH₂-cyclopentyl, cyclohexyl, -CH₂-cyclohexyl, cyclohexenylethyl, cyclohexanylethyl, norborbyl moieties and the like, which again, may bear one or more substituents.

[0024] In general, the term “cycloheteroaliphatic”, as used herein, refers to a cyclic heteroaliphatic moiety, wherein the term *heteroaliphatic* is as defined above. A cycloheteroaliphatic moiety may be substituted or unsubstituted and saturated or unsaturated. Substituents include, but are not limited to, any of the previously mentioned substituents, *i.e.*, the substituents recited for aliphatic moieties, or for other moieties as disclosed herein, resulting in the formation of a stable compound. The term “cycloheteroaliphatic” encompasses “heterocycloalkyl”, “heterocycle” or “heterocyclic” moieties, as defined herein.

[0025] Additionally, it will be appreciated that any of the cycloaliphatic or cycloheteroaliphatic moieties described above and herein may comprise an aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl moiety fused thereto. Additional examples of generally applicable substituents are illustrated by the specific embodiments shown in the Examples that are described herein.

[0026] The terms “halo” and “halogen” as used herein refer to an atom selected from fluorine, chlorine, bromine and iodine.

[0027] The term “haloalkyl” denotes an alkyl group, as defined above, having one, two, or three halogen atoms attached thereto and is exemplified by such groups as chloromethyl, bromoethyl, trifluoromethyl, and the like.

[0028] The term “acyl”, as used herein, refers to a group having the general formula -C(=O)R, where R is an aliphatic, heteroaliphatic, heterocycle, aromatic or heteroaromatic moiety, as defined herein.

[0029] The term “heterocycloalkyl”, “heterocycle” or “heterocyclic”, as used herein, refers to compounds which combine the properties of heteroaliphatic and cyclic compounds and include, but are not limited to, saturated and unsaturated mono- or polycyclic cyclic ring systems having 5-16 atoms wherein at least one ring atom is a heteroatom selected from O, S and N (wherein the nitrogen and sulfur heteroatoms may be optionally be oxidized), wherein the ring systems are optionally

substituted with one or more functional groups, as defined herein. In certain embodiments, the term "heterocycloalkyl", "heterocycle" or "heterocyclic" refers to a non-aromatic 5-, 6- or 7- membered ring or a polycyclic group wherein at least one ring atom is a heteroatom selected from O, S and N (wherein the nitrogen and sulfur heteroatoms may be optionally be oxidized), including, but not limited to, a bi- or tri-cyclic group, comprising fused six-membered rings having between one and three heteroatoms independently selected from oxygen, sulfur and nitrogen, wherein (i) each 5-membered ring has 0 to 2 double bonds, each 6-membered ring has 0 to 2 double bonds and each 7-membered ring has 0 to 3 double bonds, (ii) the nitrogen and sulfur heteroatoms may be optionally be oxidized, (iii) the nitrogen heteroatom may optionally be quaternized, and (iv) any of the above heterocyclic rings may be fused to an aryl or heteroaryl ring. Representative heterocycles include, but are not limited to, heterocycles such as furanyl, thiofuranyl, pyranal, pyrrolyl, thienyl, pyrrolidinyl, pyrazolinyl, pyrazolidinyl, imidazolinyl, imidazolidinyl, piperidinyl, piperazinyl, oxazolyl, oxazolidinyl, isooxazolyl, isoxazolidinyl, dioxazolyl, thiadiazolyl, oxadiazolyl, tetrazolyl, triazolyl, thiatriazolyl, oxatriazolyl, thiadiazolyl, oxadiazolyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, dithiazolyl, dithiazolidinyl, tetrahydrofuryl, and benzofused derivatives thereof. In certain embodiments, a "substituted heterocycle, or heterocycloalkyl or heterocyclic" group is utilized and as used herein, refers to a heterocycle, or heterocycloalkyl or heterocyclic group, as defined above, substituted by the independent replacement of one, two or three of the hydrogen atoms thereon with but are not limited to aliphatic; cycloaliphatic; heteroaliphatic; heterocyclic; aromatic; heteroaromatic; aryl; heteroaryl; alkylaryl; heteroalkylaryl; alkylheteroaryl; heteroalkylheteroaryl; alkoxy; aryloxy; heteroalkoxy; heteroaryloxy; alkylthio; arylthio; heteroalkylthio; heteroarylthio; F; Cl; Br; I; - OH; -NO₂; -CN; -CF₃; -CH₂CF₃; -CHCl₂; -CH₂OH; -CH₂CH₂OH; -CH₂NH₂; -CH₂SO₂CH₃; -C(O)R_x; -CO₂(R_x); -CON(R_x)₂; -OC(O)R_x; -OCO₂R_x; -OCON(R_x)₂; -N(R_x)₂; -S(O)₂R_x; -NR_x(CO)R_x; -N(R_x)CO₂R_x; -N(R_x)S(O)₂R_x; -N(R_x)C(=O)N(R_x)₂; -S(O)₂N(R_x)₂; wherein each occurrence of R_x independently includes, but is not limited to, aliphatic, cycloaliphatic, heteroaliphatic, heterocyclic, aromatic, heteroaromatic, aryl, heteroaryl, alkylaryl, alkylheteroaryl, heteroalkylaryl or

heteroalkylheteroaryl, wherein any of the aliphatic, cycloaliphatic, heteroaliphatic, heterocyclic, alkylaryl, or alkylheteroaryl substituents described above and herein may be substituted or unsubstituted, branched or unbranched, saturated or unsaturated, and wherein any of the aromatic, heteroaromatic, aryl or heteroaryl substituents described above and herein may be substituted or unsubstituted. Additional examples or generally applicable substituents are illustrated by the specific embodiments shown in the Examples, which are described herein.

[0030] As used herein, the terms "aliphatic", "heteroaliphatic", "alkyl", "alkenyl", "alkynyl", "heteroalkyl", "heteroalkenyl", "heteroalkynyl", and the like encompass substituted and unsubstituted, saturated and unsaturated, and linear and branched groups. Similarly, the terms "alicyclic", "heterocyclic", "heterocycloalkyl", "heterocycle" and the like encompass substituted and unsubstituted, and saturated and unsaturated groups. Additionally, the terms "cycloalkyl", "cycloalkenyl", "cycloalkynyl", "heterocycloalkyl", "heterocycloalkenyl", "heterocycloalkynyl", "aromatic", "heteroaromatic", "aryl", "heteroaryl" and the like encompass both substituted and unsubstituted groups.

[0031] As used herein, the term "isolated", when applied to the compounds of the present invention, refers to such compounds that are (i) separated from at least some components with which they are associated in nature or when they are made and/or (ii) produced, prepared or manufactured by the hand of man.

[0032] The phrase, "pharmaceutically acceptable derivative", as used herein, denotes any pharmaceutically acceptable salt, ester, or salt of such ester, of such compound, or any other adduct or derivative which, upon administration to a patient, is capable of providing (directly or indirectly) a compound as otherwise described herein, or a metabolite or residue thereof. Pharmaceutically acceptable derivatives thus include among others pro-drugs. A pro-drug is a derivative of a compound, usually with significantly reduced pharmacological activity, which contains an additional moiety that is susceptible to removal *in vivo* yielding the parent molecule as the pharmacologically active species. An example of a pro-drug is an ester which is cleaved *in vivo* to yield a compound of interest. Pro-drugs of a variety of compounds, and materials and methods for derivatizing the parent compounds to create the pro-drugs, are known and may be adapted to the present invention.

Certain exemplary pharmaceutical compositions and pharmaceutically acceptable derivatives will be discussed in more detail herein below.

[0033] The term “treating”, as used herein generally means that the compounds of the invention can be used in humans or animals with at least a tentative diagnosis of disease. The compounds of the invention will delay or slow the progression of the disease thereby giving the individual a more useful life span.

[0034] The term “preventing” as used herein means that the compounds of the present invention are useful when administered to a patient who has not been diagnosed as possibly having the disease at the time of administration, but who would normally be expected to develop the disease or be at increased risk for the disease. The compounds of the invention will slow the development of disease symptoms, delay the onset of disease, or prevent the individual from developing the disease at all. Preventing also includes administration of the compounds of the invention to those individuals thought to be predisposed to the disease due to age, familial history, genetic or chromosomal abnormalities, and/or due to the presence of one or more biological markers for the disease, such as a known genetic mutation of APP or APP cleavage products in brain tissues or fluids.

[0035] As used herein the term “*biological sample*” includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from an animal (*e.g.*, mammal) or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof. For example, the term “biological sample” refers to any solid or fluid sample obtained from, excreted by or secreted by any living organism, including single-celled micro-organisms (such as bacteria and yeasts) and multicellular organisms (such as plants and animals, for instance a vertebrate or a mammal, and in particular a healthy or apparently healthy human subject or a human patient affected by a condition or disease to be diagnosed or investigated). The biological sample can be in any form, including a solid material such as a tissue, cells, a cell pellet, a cell extract, cell homogenates, or cell fractions; or a biopsy, or a biological fluid. The biological fluid may be obtained from any site (*e.g.* blood, saliva (or a mouth wash containing buccal cells), tears, plasma, serum, urine, bile, cerebrospinal fluid, amniotic fluid, peritoneal fluid, and pleural fluid, or cells therefrom, aqueous or vitreous humor, or any bodily secretion), a transudate, an

exudate (*e.g.* fluid obtained from an abscess or any other site of infection or inflammation), or fluid obtained from a joint (*e.g.* a normal joint or a joint affected by disease such as rheumatoid arthritis, osteoarthritis, gout or septic arthritis). The biological sample can be obtained from any organ or tissue (including a biopsy or autopsy specimen) or may comprise cells (whether primary cells or cultured cells) or medium conditioned by any cell, tissue or organ. Biological samples may also include sections of tissues such as frozen sections taken for histological purposes. Biological samples also include mixtures of biological molecules including proteins, lipids, carbohydrates and nucleic acids generated by partial or complete fractionation of cell or tissue homogenates. Although the sample is preferably taken from a human subject, biological samples may be from any animal, plant, bacteria, virus, yeast, etc. The term *animal*, as used herein, refers to humans as well as non-human animals, at any stage of development, including, for example, mammals, birds, reptiles, amphibians, fish, worms and single cells. Cell cultures and live tissue samples are considered to be pluralities of animals. In certain exemplary embodiments, the non-human animal is a mammal (*e.g.*, a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a sheep, cattle, a primate, or a pig). An animal may be a transgenic animal or a human clone. If desired, the biological sample may be subjected to preliminary processing, including preliminary separation techniques.

DESCRIPTION OF CERTAIN PREFERRED EMBODIMENTS OF THE INVENTION

[0036] As noted above, there has been increasing interest in recent years in the development of aspartyl protease inhibitors, particularly BACE inhibitors, as therapeutic agents for the treatment of Alzheimer's Disease and other disorders caused by the accumulation of β -amyloid plaques. It has been generally accepted by the scientific community that potential new leads for aspartyl inhibitor small molecules must comprise an aspartate binding hydroxyl group to retain aspartyl protease inhibitory activity (See for example, Ajay *et al.*, "Designing Libraries with CNS Activity", *J. Med. Chem.*, 42:4942-4951, 1999, see especially the paragraph bridging columns 1 and 2 on page 4942 of this article). The present invention demonstrates that amino analogs (*i.e.*, where the hydroxyl group has been replaced with an amino group) are unexpectedly equally promising (if not superior) aspartyl

protease inhibitors as their hydroxy-containing counterparts. In fact, the present invention shows that certain amino-containing inventive compounds possess several superior biological properties over their hydroxy-containing counterparts (*e.g.*, increased potency in cells, increased selectivity for the BACE enzyme, and/or superior ADME properties).


[0037] Thus, the present invention provides novel amino-containing compounds capable of inhibiting the activity of BACE. More generally, the compounds of the invention are inhibitors of proteases, and more specifically inhibitors of aspartyl proteases. In certain embodiments of special interest, the inventive compounds are useful for the treatment or prevention of disorders characterized by β amyloid deposits or plaques. In certain exemplary embodiments, the compounds are useful for the treatment of Alzheimer's Disease.

[0038] Compounds of this invention include those generally set forth above and described specifically herein, and are illustrated in part by the various classes, subgenera and species disclosed herein.

[0039] Additionally, the present invention provides pharmaceutically acceptable derivatives of the inventive compounds, and methods of treating a subject using these compounds, pharmaceutical compositions thereof, or either of these in combination with one or more additional therapeutic agents.

[0040] *1) General Description of Compounds of the Invention*

[0041] In certain embodiments, the compounds of the invention include compounds of the general formula (I) as further defined below:



and pharmaceutically acceptable derivatives thereof;

wherein R' is hydrogen or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R' , taken together with R^2 or a substituent present on R^1 , may form a cycloheteroaliphatic moiety;

R^1 is an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^1 , taken together with R' , may form a cycloheteroaliphatic moiety;

X^1 is $-C(=O)-$, $-S(=O)-$, $-C(=NH)-$, $-C(=S)-$, $-NC(=O)-$, $-NC(=S)-$, $-N-C(=N-C\equiv N)-$, $-NS(O_2)-$, $-CHR^{X1A}-$, $-SO_2-$, $-COO-$, $-C(=O)C(R^{X1A})_2-$, or $-SC(=O)-$ wherein each occurrence of R^{X1A} is independently hydrogen, or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety;

R^2 is an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^2 , taken together with R^1 , may form a cycloheteroaliphatic moiety;

R^3 is hydrogen, halogen, or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety;

$R^{3'}$ is hydrogen, halogen, or lower alkyl;

R^4 is hydrogen, an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^4 , taken together with a substituent present on X^2 or X^3 , may form a cycloaliphatic, cycloheteroaliphatic, aromatic, or heteroaromatic moiety;

X^2 is absent, $-NR^{X2A}-$, $-(CHR^{X2A})_j-$, $-NR^{X2A}Y-$, $-(CHR^{X2A})_jY-$ or $-N(R^{X2A})CH(R^{X2A})Y-$ wherein each occurrence of R^{X2A} is independently hydrogen or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety; each occurrence

of Y is independently $\left(\begin{array}{c} \text{O} \\ \parallel \\ \text{---} \text{N} \text{---} \text{CH} \text{---} \text{R}^{X2B} \\ | \\ \text{H} \end{array} \right)_t$ wherein, for each independent occurrence of t , R^{X2B} is hydrogen, or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^{X2A} or one occurrence of R^{X2B} taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aromatic or heteroaromatic moiety, and wherein each occurrence of j and t is independently an integer from 1 to 4; and

X^3 is absent, $-NHCO-$, $-NHSO_2-$, $-NHCONH-$, $-NHCOO-$, $-CH_2NH-$, $-C(=O)-$, $-S(=O)-$, $-C(=NH)-$, $-C(=S)-$, $-NC(=S)N-$, $-N-C(=N-C\equiv N)N-$, $-NS(O_2)N-$, $-SO_2-$, $-C(=O)NR^{X3A}-$, $-C(=S)NR^{X3A}-$, $-COO-$, $-(CHR^{X3A})_k-$, $-O-$, $-CH_2NR^{X3A}-$, or $-NR^{X3A}-$, wherein each occurrence of R^{X3A} is independently hydrogen, an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^{X3A} taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aromatic or heteroaromatic moiety, and k is an integer from 1 to 3.

[0042] In certain embodiments of compounds described directly above and compounds as described in certain classes and subclasses herein, one or more of the following groups do not occur simultaneously as defined:

(i) R^1 , R^3 and $R^{3'}$ are each hydrogen; R^2 is alkyl, cycloalkylalkyl or aralkyl; $-X^2-X^3-R^4$ together represents $-\text{CHR}^c\text{C}(=\text{O})\text{NHCH}(\text{R}^w)\text{C}(=\text{O})\text{NR}^x\text{R}^y$, wherein R^c is hydrogen or alkyl, R^w is alkyl, and one of R^x or R^y represents hydrogen and the other represents hydrogen, alkyl, aryl, aralkyl, 1-alkoxycarbonyl-2-phenylethyl, 1-alkoxycarbonyl-2-(imidazol-4-yl)ethyl, 2-(imidazol-1-yl)ethyl, indanyl, heterocyclyl-alkyl, carboxyalkyl, alkoxycarbonylalkyl, aryloxycarbonylalkyl, aralkoxycarbonylalkyl or a group of the formula $-\text{A}-\text{N}(\text{R}^a)(\text{R}^b)$ in which A represents alkylene and R^a and R^b each represents alkyl or R^a and R^b together represent a pentamethylene group in which one methylene group can be replaced by NH, N-alkyl, N-alkanoyl, N-aralkoxycarbonyl, O, S, SO, or SO_2 ; or R^x and R^y together with the nitrogen atom to which they are attached represent a 1,2,3,4-tetrahydroisoquinoline ring; and $-X^1-R^1$ together represents an alkoxycarbonyl, aralkoxycarbonyl, alkanoyl, aralkanoyl, aroyl, cycloalkylcarbonyl, heterocyclylcarbonyl, heterocyclyl-alkanoyl, 6-(dibenzylcarbamoyle)-4-oxohexanoyl moiety or an acyl group of an α -amino acid in which the amino group is substituted by an alkoxycarbonyl, aralkoxycarbonyl, diaralkylcarbamoyle, diaralkylalkanoyl, or aralkanoyl moiety;

wherein the term "aroyl" refers to an acyl group derived from an arylcarboxylic acid such as benzoyl, 1-naphthoyl, 2-naphthoyl, etc., and the term "aralkanoyl" refers to an acyl group derived from an aryl-substituted alkanecarboxylic acid;

whereby the term "aryl" alone or in each of the aralkyl, aryloxycarbonylalkyl, aralkoxycarbonylalkyl or N-aralkoxycarbonyl moieties refers to a phenyl or naphthyl group optionally substituted with one or more substituents selected from alkyl, hydroxy, alkoxy and halogen;

(ii) R^1 , R^3 and $R^{3'}$ are each hydrogen; X^1 is $-\text{C}(=\text{O})-$, $-\text{SO}_2-$, $\text{N}(\text{R}^x)\text{SO}_2$, $\text{N}(\text{R}^x)\text{C}(=\text{O})$ or $\text{SC}(=\text{O})$, wherein R^x is hydrogen, C_{1-5} alkyl or joined together with R^1 either directly to form a 5-7 membered heterocycle such as pyrrolidinyl or piperidinyl, or through a heteroatom selected from N, O and S, to form a 6-

membered heterocycle with the nitrogen to which they are attached such as morpholinyl, piperazyl, or N-C₁₋₃alkyl-piperazyl; R¹ is a substituted or unsubstituted C₁₋₆alkyl, a 5-6 membered heterocycle or a 6-10 carbon atoms aryl moiety substituted with C₁₋₄alkyl, C₁₋₃alkoxy, hydroxy, halogen, N(R^a)₂, C(=O)OR^a, -C(=O)N(R^a)₂, -SO₂N(R^a)₂, CH₂N(R^a)₂, N(R^a)C(=O)R^a or N(R^a)SO₂R^a; R² is OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c; and -X²-X³-R⁴ together represent -CH(R^d)C(=O)NHCH(R^e)C(=O)-Y-[CR^fR^g]_mR^g, wherein m is an integer from 0 to 5, Y is O or NH, R^d and R^e are independently hydrogen, OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c wherein n is an integer from 0 to 5, R^a is hydrogen or C₁₋₄alkyl, R^b is hydrogen, hydroxy or C₁₋₄alkyl and R^c is hydrogen, substituted or unsubstituted aryl, 5- or 6-membered heterocycle, C₁₋₆alkyl or C₁₋₆alkenyl, C₃₋₇cycloalkyl, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring, benzofuryl, indolyl, azabicyclo C₇₋₁₁cycloalkyl or benzopiperidinyl; R^f is hydrogen, substituted or unsubstituted C₁₋₆alkyl or (CH₂CH₂O)_pCH₃ or (CH₂CH₂O)_pH wherein p is an integer from 0 to 5, and R^g is hydrogen, or substituted or unsubstituted aryl, heterocycle, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring;

(iii) R¹, R³ and R^{3'} are each hydrogen; X¹ is -C(=O)-, -SO₂-, N(R^x)SO₂, N(R^x)C(=O) or SC(=O), wherein R^x is hydrogen, C₁₋₅alkyl or joined together with R¹ either directly to form a 5-7 membered heterocycle such as pyrrolidinyl or piperidinyl, or through a heteroatom selected from N, O and S, to form a 6-membered heterocycle with the nitrogen to which they are attached such as morpholinyl, piperazyl, or N-C₁₋₃alkyl-piperazyl; R¹ is a substituted or unsubstituted C₁₋₆alkyl, a 5-6 membered heterocycle or a 6-10 carbon atoms aryl moiety substituted with C₁₋₄alkyl, C₁₋₃alkoxy, hydroxy, halogen, N(R^a)₂, C(=O)OR^a, -C(=O)N(R^a)₂, -SO₂N(R^a)₂, CH₂N(R^a)₂, N(R^a)C(=O)R^a or N(R^a)SO₂R^a; R² is OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c; and -X²-X³-R⁴ together represent -CH(R^d)C(=O)-Y-[CR^fR^g]_mR^g, wherein m is an integer from 0 to 5, Y is O or NH, R^d is hydrogen, OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c; wherein n is an integer from 0 to 5, R^a is hydrogen or C₁₋₄alkyl, R^b is hydrogen, hydroxy or C₁₋₄alkyl and R^c is hydrogen, substituted or unsubstituted aryl, 5- or 6-membered heterocycle, C₁₋₆alkyl or C₁₋₆alkenyl, C₃₋₇cycloalkyl, 5- to 7-membered carbocyclic or 7- to 10-

membered bicyclic carbocyclic ring, benzofuryl, indolyl, azabicyclo C₇₋₁₁cycloalkyl or benzopiperidinyl; R^f is hydrogen, substituted or unsubstituted C₁₋₆alkyl or (CH₂CH₂O)_pCH₃ or (CH₂CH₂O)_pH wherein p is an integer from 0 to 5, and R^g is hydrogen, or substituted or unsubstituted aryl, heterocycle, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring;

(iv) R' and R^{3'} are each hydrogen; X¹ is -C(=O)-, -SO₂-, N(R^x)SO₂, N(R^x)C(=O) or SC(=O), wherein R^x is hydrogen, C₁₋₅alkyl or joined together with R¹ either directly to form a 5-7 membered heterocycle such as pyrrolidinyl or piperidinyl, or through a heteroatom selected from N, O and S, to form a 6-membered heterocycle with the nitrogen to which they are attached such as morpholinyl, piperazyl, or N-C₁₋₃alkyl-piperazyl; R¹ is a substituted or unsubstituted C₁₋₆alkyl, a 5-6 membered heterocycle or a 6-10 carbon atoms aryl moiety substituted with C₁₋₄alkyl, C₁₋₃alkoxy, hydroxy, halogen, N(R^a)₂, C(=O)OR^a, -C(=O)N(R^a)₂, -SO₂N(R^a)₂, CH₂N(R^a)₂, N(R^a)C(=O)R^a or N(R^a)SO₂R^a; R² is OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c; R³ is hydrogen, OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c; and -X²-X³-R⁴ together represent -CH(R^d)C(=O)NHCH(R^e)C(=O)-Y-[CR^fR^g]_mR^g, wherein m is an integer from 0 to 5, Y is O or NH, R^d and R^e are independently hydrogen, OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c wherein n is an integer from 0 to 5, R^a is hydrogen or C₁₋₄alkyl, R^b is hydrogen, hydroxy or C₁₋₄alkyl and R^c is hydrogen, substituted or unsubstituted aryl, 5- or 6-membered heterocycle, C₁₋₆alkyl or C₁₋₆alkenyl, C₃₋₇cycloalkyl, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring, benzofuryl, indolyl, azabicyclo C₇₋₁₁cycloalkyl or benzopiperidinyl; R^f is hydrogen, substituted or unsubstituted C₁₋₆alkyl or (CH₂CH₂O)_pCH₃ or (CH₂CH₂O)_pH wherein p is an integer from 0 to 5, and R^g is hydrogen, or substituted or unsubstituted aryl, heterocycle, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring;


(v) R' and R^{3'} are each hydrogen; X¹ is -C(=O)-, -SO₂-, N(R^x)SO₂, N(R^x)C(=O) or SC(=O), wherein R^x is hydrogen, C₁₋₅alkyl or joined together with R¹ either directly to form a 5-7 membered heterocycle such as pyrrolidinyl or piperidinyl, or through a heteroatom selected from N, O and S, to form a 6-membered heterocycle with the nitrogen to which they are attached such as morpholinyl, piperazyl, or N-C₁₋₃alkyl-piperazyl; R¹ is a substituted or unsubstituted

C₁₋₆alkyl, a 5-6 membered heterocycle or a 6-10 carbon atoms aryl moiety substituted with C₁₋₄alkyl, C₁₋₃alkoxy, hydroxy, halogen, N(R^a)₂, C(=O)OR^a, -C(=O)N(R^a)₂, -SO₂N(R^a)₂, CH₂N(R^a)₂, N(R^a)C(=O)R^a or N(R^a)SO₂R^a; R² is OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c; R³ is hydrogen, OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c; and -X²-X³-R⁴ together represent -CH(R^d)C(=O)-Y-[CR^fR^g]_mR^g, wherein m is an integer from 0 to 5, Y is O or NH, R^d is hydrogen, OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c; wherein n is an integer from 0 to 5, R^a is hydrogen or C₁₋₄alkyl, R^b is hydrogen, hydroxy or C₁₋₄alkyl and R^c is hydrogen, substituted or unsubstituted aryl, 5- or 6-membered heterocycle, C₁₋₆alkyl or C₁₋₆alkenyl, C₃₋₇cycloalkyl, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring, benzofuryl, indolyl, azabicyclo C₇₋₁₁cycloalkyl or benzopiperidinyl; R^f is hydrogen, substituted or unsubstituted C₁₋₆alkyl or (CH₂CH₂O)_pCH₃ or (CH₂CH₂O)_pH wherein p is an integer from 0 to 5, and R^g is hydrogen, or substituted or unsubstituted aryl, heterocycle, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring; and


(vi) R¹, R³ and R^{3'} are each hydrogen; R² is C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₇cycloalkyl, aryl, heteroaryl, T-C₁₋₆alkyl, T-C₂₋₆alkenyl, wherein T is aryl, heteroaryl or C₃₋₇cycloalkyl; R¹-X¹ together represent W wherein W is R^x, R^xCO, R^xOCO, R^xOCH(R^y)CO, R^xNHCH(R^y)CO, R^xSCH(R^y)CO, R^xSO₂, R^xSO or an amino acid with a blocked or unblocked amino terminus, wherein R^x and R^y are each independently hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, aryl, heteroaryl, T-C₁₋₆alkyl or T-(CH₂)_nCH(T)(CH₂)_n wherein n is an integer from 1 to 4; and -X²-X³-R⁴ together represent -CH(R^a)C(=X)CHR^bR^c, wherein X is (OH,H) or O; R^a is hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₇cycloalkyl, aryl, heteroaryl, T-C₁₋₆alkyl or T-C₂₋₆alkenyl; R^b is hydrogen or OH; and R^c is Y, (CHR^w)_n-Y or =CR^z(CHR^w)_n-Y, wherein Y is hydrogen, OH, -NR^wR^q, aryl, heteroaryl or CO-Z, n is an integer from 1 to 4, Z is OH, -NR^wR^q, OR^w or an amino acid with a blocked or unblocked carboxy terminus, R^q is H, C₁₋₆alkyl or arylC₁₋₆alkyl, and R^z and R^w are each independently hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, aryl, heteroaryl, T-C₁₋₆alkyl or T-C₂₋₆alkenyl.

[0043] In certain embodiments, the present invention defines particular classes of compounds which are of special interest. For example, one class of compounds


of special interest includes those compounds of formula (I) having the stereochemistry shown in Formula (I^A):




[0044] Another class of compounds of special interest includes those compounds of formula (I^A) wherein, X² is absent and the compound has the Formula (I^B):




[0045] Another class of compounds of special interest includes those compounds of formula (I^A) wherein X³ is absent and the compound has the Formula (I^C):



[0046] Another class of compounds of special interest includes those compounds of formula (I^A) wherein X² and X³ are each absent and the compound has the Formula (I^D):




[0047] Another class of compounds of special interest includes those compounds of formula (I^A) wherein R^{3'} is hydrogen, X² is CHMe and the compound has the Formula (I^E):




[0048] Another class of compounds of special interest includes those compounds of formula (I^A) wherein R^{3'} is hydrogen, X³ is absent, X² is -CH(Me)Y-


where Y is $\left(\begin{array}{c} \text{O} \\ \parallel \\ \text{---} \text{N} \text{---} \text{CH} \text{---} \text{R}^{\text{X2B}} \end{array} \right)_2$ and the compound has the Formula (I^F):




[0049] Another class of compounds of special interest includes those compounds of formula (I) wherein R' and R^{3'} are each hydrogen, X² is CHMe and X³ is -C(=O)NH-, and the compound has the Formula (II):




[0050] Another class of compounds of special interest includes those compounds of formula (I) wherein X¹ is -C(=O)-, R' is hydrogen, X² and X³ are each absent, and the compound has the Formula (III):



[0051] Another class of compounds of special interest includes those compounds of formula (I) wherein X¹ is -C(=O)-, R' and R^{3'} are each hydrogen, X² is -CHMe, and the compound has the Formula (IV):




[0052] Another class of compounds of special interest includes those compounds of formula (I) wherein R' and R^{3'} are each hydrogen, X² is -NR^{X2A}-, and the compound has the Formula (V):




(V)

[0053] Another class of compounds of special interest includes those compounds of formula (V) above having the stereochemistry as shown in Formula (V^A):


(V^A)

[0054] Another class of compounds of special interest includes those compounds of formula (I) wherein R' and R^{3'} are each hydrogen, X¹ is -C(=O)-, X² is -NH-, and the compound has the Formula (VI):




(VI)

[0055] Another class of compounds of special interest includes those compounds of Formula (I) wherein X¹ is -C(=O)- and the compound has the structure as shown in Formula (VII):




(VII)

[0056] Another class of compounds of special interest includes those compounds of formula (I) wherein X¹ is CHR^{X1A} and the compound has the structure as shown in Formula (VIII):




(VIII)


[0057] Another class of compounds of special interest includes those compounds of formula (I) wherein X^1 is CH_2 , X^2 is CHR^{X2A} , X^3 is $-\text{C}(=\text{O})\text{NR}^{X3A}$, and the compound has the Formula (IX):




[0058] Another class of compounds of special interest includes those compounds of formula (IX) wherein R^{X2A} is methyl and the compound has the Formula (IX^A):



[0059] Another class of compounds of special interest includes those compounds of formula (I) wherein X^1 is $-\text{C}(=\text{O})-$, X^2 is CHR^{X2A} , X^3 is $-\text{CH}_2\text{NR}^{X3A}$, and the compound has the Formula (X):



[0060] Another class of compounds of special interest includes those compounds of formula (X) wherein R^{X2A} is methyl and the compound has the Formula (X^A):



[0061] In certain other exemplary embodiments, for each of the classes of compounds as described above, R^3 is hydrogen; R^2 is substituted or unsubstituted lower alkyl, lower alkylamino, lower heteroaryl, $-(\text{CH}_2)\text{cycloalkyl}$, $-(\text{CH}_2)\text{heterocycloalkyl}$, $-(\text{CH}_2)\text{aryl}$, $-(\text{CH}_2)\text{heteroaryl}$, optionally substituted with one or more occurrences of R^{2A} , wherein R^{2A} is hydrogen, alkyl, heteroalkyl, aryl,

heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -OR^{2B}, -SR^{2B}, -N(R^{2B})₂, -SO₂N(R^{2B})₂, -C(=O)N(R^{2B})₂, halogen, -CN, -NO₂, -C(=O)OR^{2B}, -N(R^{2B})C(=O)R^{2C}, wherein each occurrence of R^{2B} and R^{2C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl; R¹ is alkyl, alkenyl, heteroalkyl, heteroalkenyl, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl, wherein R¹ is optionally substituted with one or more occurrences of R^{1A}, wherein R^{1A} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -OR^{1B}, -SR^{1B}, -N(R^{1B})₂, -SO₂N(R^{1B})₂, -C(=O)N(R^{1B})₂, halogen, -CN, -NO₂, -C(=O)OR^{1B}, -N(R^{1B})C(=O)R^{1C}, or R^{1B} and R^{1C}, taken together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclic moiety; wherein each occurrence of R^{1B} and R^{1C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl; and R⁴ is alkyl, alkenyl, heteroalkyl, heteroalkenyl, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl, optionally substituted with one or more occurrences of R^{4A}, wherein R^{4A} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -OR^{4B}, -SR^{4B}, -N(R^{4B})₂, -SO₂N(R^{4B})₂, -C(=O)N(R^{4B})₂, halogen, -CN, -NO₂, -C(=O)OR^{4B}, -N(R^{4B})C(=O)R^{4C}, wherein each occurrence of R^{4B} and R^{4C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl.


[0062] A number of important subclasses of each of the foregoing classes deserve separate mention; these subclasses include subclasses of the foregoing classes in which:

[0063] i) R¹ is substituted or unsubstituted, linear or branched, cyclic or acyclic alkyl or alkenyl;

[0064] ii) R¹ is substituted or unsubstituted aryl, heteroaryl, -(alkyl)aryl, -


(alkyl)heteroaryl or heterocyclic;

[0065] iii) R¹ is one of:




wherein R^{1A} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, $-\text{OR}^{1B}$, $-\text{SR}^{1B}$, $-\text{N}(\text{R}^{1B})_2$, $-\text{SO}_2\text{N}(\text{R}^{1B})_2$, $-\text{C}(=\text{O})\text{N}(\text{R}^{1B})_2$, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OR}^{1B}$, $\text{N}(\text{R}^{1B})\text{C}(=\text{O})\text{R}^{1C}$ or $-\text{N}(\text{R}^{1B})\text{SO}_2\text{R}^{1C}$; wherein each occurrence of R^{1B} and R^{1C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$; or R^{1B} and R^{1C} , taken together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclic moiety; R^{1D} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, acyl or a nitrogen protecting group; wherein n and p are each independently integers from 0 to 3 and r is an integer from 1 to 6; whereby each of the foregoing alkyl and heteroalkyl moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, and each of the foregoing aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ and $-(\text{alkyl})\text{heteroaryl}$ moieties may be substituted or unsubstituted;

[0066] iv) R^1 is one of:




wherein n , p , R^{1A} and R^{1D} are as defined in iii) above;

[0067] v) R^1 is one of:




wherein n , p , R^{1A} and R^{1B} are as defined in iii) above; R^{1D} is hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(alkyl)aryl$; $-(alkyl)heteroaryl$ or acyl; R^{1E} and R^{1F} are each independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(alkyl)aryl$, $-(alkyl)heteroaryl$, $-C(=O)R^{1C}$ or $-SO_2R^{1C}$, where R^{1C} is hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(alkyl)aryl$ or $-(alkyl)heteroaryl$; or R^{1E} and R^{1F} taken together form a 5-8 membered heterocyclic ring; or R^{1E} and one occurrence of R^{1A} , taken together, form a substituted or unsubstituted, saturated or unsaturated heterocyclic ring;

[0068] vi) R^1 is:




wherein n is 1-2; p is 0-1; R^{1A} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ or $-(\text{alkyl})\text{heteroaryl}$, $-\text{OR}^{1B}$, $-\text{SR}^{1B}$, $-\text{N}(\text{R}^{1B})_2$, $-\text{SO}_2\text{N}(\text{R}^{1B})_2$, $-\text{C}(=\text{O})\text{N}(\text{R}^{1B})_2$, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OR}^{1B}$, $-\text{N}(\text{R}^{1B})\text{C}(=\text{O})\text{R}^{1C}$ or $-\text{N}(\text{R}^{1B})\text{SO}_2\text{R}^{1C}$, wherein each occurrence of R^{1B} and R^{1C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ or $-(\text{alkyl})\text{heteroaryl}$; or R^{1B} and R^{1C} , taken together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclic moiety; R^{1D} and R^{1E} are each independently hydrogen or lower alkyl; or R^{1D} and R^{1E} taken together form a 5-8 membered heterocyclic ring; or R^{1E} and one occurrence of R^{1A} , taken together, form a substituted or unsubstituted, saturated or unsaturated heterocyclic ring;

[0069] vii) R^1 is:




wherein a is 0 or 1; n is from 0-4; and R^{1A} and R^{1D} are as defined in vii) above;

[0070] viii) R^1 is:




wherein R^{1A} is hydrogen, halogen, $-\text{CN}$, $-\text{NO}_2$, lower alkyl, lower heteroalkyl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, $-\text{OR}^{1B}$ or $-\text{N}(\text{R}^{1B})\text{SO}_2\text{R}^{1C}$; wherein each occurrence of R^{1B} and R^{1C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$; whereby each of the foregoing alkyl and heteroalkyl moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, and each of the foregoing aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ and $-(\text{alkyl})\text{heteroaryl}$ moieties may be substituted or unsubstituted;

[0071] ix) R^1 is:




wherein R^{1A} is as defined in viii) above; and R^{1D} is hydrogen or lower alkyl;

[0072] x) R^1 is:




wherein n is 1-2; p is 0-1; R^{1A} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl, $-OR^{1B}$, $-SR^{1B}$, $-N(R^{1B})_2$, $-SO_2N(R^{1B})_2$, $-C(=O)N(R^{1B})_2$, halogen, $-CN$, $-NO_2$, $-C(=O)OR^{1B}$, $-N(R^{1B})C(=O)R^{1C}$ or $-N(R^{1B})SO_2R^{1C}$, wherein each occurrence of R^{1B} and R^{1C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl; or R^{1B} and R^{1C} , taken together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclic moiety; R^{1D} and R^{1E} are each independently hydrogen or lower alkyl; or R^{1D} and R^{1E} taken together form a 5-8 membered heterocyclic ring; or R^{1E} and one occurrence of R^{1A} , taken together, form a substituted or unsubstituted, saturated or unsaturated heterocyclic ring;

[0073] xi) R^1 is:



wherein R^{1A} is hydrogen, halogen, $-CN$, $-NO_2$, lower alkyl, lower heteroalkyl, -(alkyl)aryl, -(alkyl)heteroaryl, $-OR^{1B}$ or $-N(R^{1B})SO_2R^{1C}$; wherein each occurrence of R^{1B} and R^{1C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl; whereby each of the foregoing alkyl and heteroalkyl moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, and each of the foregoing aryl, heteroaryl, -(alkyl)aryl and -(alkyl)heteroaryl moieties may be substituted or unsubstituted;

[0074] xii) R^1 is:



wherein R^{1A} is as defined in x) above; and R^{1D} is hydrogen or lower alkyl;


[0075] xiii) compounds of subsets vi)-xii) wherein R^{1A} is methyl, methoxy or halide;

[0076] xiv) compounds of subsets vi)-xii) wherein R^{1A} is methyl, methoxy or F;

[0077] xv) compounds of subsets vi)-xii) wherein R^{1A} is methyl;

[0078] xvi) compounds of subsets viii) and xii) wherein R^{1B} is hydrogen, methyl or ethyl;

[0079] xvii) R^1 is:




wherein R^{1A} and R^{1B} are each independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl;

[0080] xviii) compounds of subset xvii) wherein R^{1A} and R^{1B} are each independently cyclic or acyclic lower alkyl;

[0081] xix) compounds of subset xvii) wherein R^{1A} and R^{1B} are each independently methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, sec-butyl, isopentyl or cyclopropyl;

[0082] xx) R^1 is:




wherein R^{1A} is hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl; R^{1E} and R^{1F} are each independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -C(=O) R^{1C} or -SO₂ R^{1C} , where R^{1C} is hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl; or R^{1E} and R^{1F} taken together form a 5-8 membered heterocyclic ring;


[0083] xxi) compounds of subset xx) wherein R^{1A} is substituted or unsubstituted, linear or branched, cyclic or acyclic lower alkyl; R^{1E} is lower alkyl; and R^{1F} is $-\text{SO}_2R^{1D}$ wherein R^{1D} is lower alkyl;

[0084] xxii) compounds of subset xx) wherein R^{1A} is substituted or unsubstituted, linear or branched, cyclic or acyclic lower alkyl; R^{1E} and R^{1F} , taken together with the nitrogen atom to which they are attached, form a 5- or 6-membered cyclic sulfonamide moiety;


[0085] xxiii) R^1 is one of:



[0086] xxiv) R^1 is one of:



[0087] xxv) R^1 is one of:




[0088] xxvi) R^2 is lower alkyl, $-\text{CH}_2\text{NR}^{2A}\text{R}^{2B}$ or $-(\text{CH}_2)\text{phenyl}$, wherein the phenyl group is optionally substituted with one or more occurrences of R^{2C} , wherein R^{2C} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ or $-(\text{alkyl})\text{heteroaryl}$, $-\text{OR}^{2D}$, $-\text{SR}^{2D}$, $-\text{N}(\text{R}^{2D})_2$, $-\text{SO}_2\text{N}(\text{R}^{2D})_2$, $-\text{C}(=\text{O})\text{N}(\text{R}^{2D})_2$, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OR}^{2D}$, $-\text{N}(\text{R}^{2D})\text{C}(=\text{O})\text{R}^{2E}$, wherein each occurrence of R^{2D} and R^{2E} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ or $-(\text{alkyl})\text{heteroaryl}$; and wherein R^{2A} and R^{2B} are each independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ or $-(\text{alkyl})\text{heteroaryl}$; whereby each of the foregoing alkyl and heteroalkyl moieties may

be linear or branched, substituted or unsubstituted, cyclic or acyclic, and each of the foregoing aryl, heteroaryl, -(alkyl)aryl and -(alkyl)heteroaryl moieties may be substituted or unsubstituted;

[0089] xxvii) R^2 is lower alkyl, $-\text{CH}_2\text{NR}^{2A}\text{R}^{2B}$ or $-(\text{CH}_2)\text{phenyl}$, wherein the phenyl group is optionally substituted with one or more occurrences of R^{2C} , wherein R^{2C} is hydrogen, alkyl, alkoxy or halogen; and wherein R^{2A} and R^{2B} are each independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl; whereby each of the foregoing alkyl and heteroalkyl moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, and each of the foregoing aryl, heteroaryl, -(alkyl)aryl and -(alkyl)heteroaryl moieties may be substituted or unsubstituted;


[0090] xxviii) R^2 is one of:



wherein R^{2A} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl, $-\text{OR}^{2B}$, $-\text{SR}^{2B}$, $-\text{N}(\text{R}^{2B})_2$, $-\text{SO}_2\text{N}(\text{R}^{2B})_2$, $-\text{C}(=\text{O})\text{N}(\text{R}^{2B})_2$, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OR}^{2B}$, $-\text{N}(\text{R}^{2B})\text{C}(=\text{O})\text{R}^{2C}$, wherein each occurrence of R^{2B} and R^{2C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl, wherein q and s are each independently integers

from 0 to 3 and u is an integer from 1 to 6; whereby each of the foregoing alkyl and heteroalkyl moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, and each of the foregoing aryl, heteroaryl, -(alkyl)aryl and -(alkyl)heteroaryl moieties may be substituted or unsubstituted;

[0091] xxix) R^2 is one of:




wherein each occurrence of R^{2A} is independently hydrogen or lower alkyl; each occurrence of X is independently a halogen; s is an integer from 0 to 3 and u is an integer from 1 to 6; whereby each of the foregoing alkyl moieties may be linear or branched, substituted or unsubstituted and cyclic or acyclic;

[0092] xxx) compounds of subset xxix) wherein X is chlorine or fluorine;


[0093] xxxi) compounds of subset xxix) wherein each occurrence of X is fluorine;

[0094] xxxii) compounds of subset xxix) wherein R^{2A} is methyl;


[0095] xxxiii) R^2 is one of:




[0096] xxxiv) R^2 is one of:



[0097] xxxv) R^2 is one of:



[0098] xxxvi) R^2 is one of:



- [0099] xxxvii) R^3 is hydrogen or halogen;
- [0100] xxxviii) R^3 is hydrogen or F;
- [0101] xxxix) R^3 is hydrogen;
- [0102] xl) $R^{3'}$ is hydrogen, methyl or halogen;
- [0103] xli) $R^{3'}$ is hydrogen, methyl or F;
- [0104] xlii) $R^{3'}$ is hydrogen;
- [0105] xliii) one of R^3 and $R^{3'}$ is halogen;
- [0106] xliv) R^3 and $R^{3'}$ are each independently hydrogen or F;
- [0107] xlv) R^3 and $R^{3'}$ are each F;
- [0108] xlvi) R^3 and $R^{3'}$ are each hydrogen;
- [0109] xlvii) X^1 is $-C(=O)-$;
- [0110] xlviii) X^1 is CHR^{X1A} and R^{X1A} is hydrogen or linear or branched substituted or unsubstituted alkyl;
- [0111] xlix) X^1 is CH_2 ;
- [0112] l) X^2 is CHR^{X2A} and R^{X2A} is hydrogen or linear or branched substituted or unsubstituted alkyl or -alkyl(aryl);
- [0113] li) X^2 is CHR^{X2A} and R^{X2A} is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, benzyl or phenethyl;
- [0114] lii) X^2 is CHR^{X2A} and R^{X2A} is methyl, ethyl, propyl, isopropyl or phenethyl;
- [0115] liii) X^2 is $CHMe$
- [0116] liv) X^2 is $-NR^{X2A}-$ and R^{X2A} is hydrogen or linear or branched substituted or unsubstituted alkyl;
- [0117] lv) X^2 is $-NR^{X2A}-$ and R^{X2A} is C1-C3 alkyl;
- [0118] lvi) X^2 is NH ;

[0119] lvii) X^2 is NHY where Y is $\left(\begin{array}{c} \text{O} \\ \parallel \\ \text{---} \text{N} \text{---} \text{CH} \text{---} \text{R}^{X2B} \\ | \\ \text{H} \end{array} \right)_2$ where each occurrence R^{X2B} is independently hydrogen, or an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, $-(\text{heteroalkyl})\text{aryl}$, or $-(\text{heteroalkyl})\text{heteroaryl}$ moiety; whereby each of the foregoing aliphatic and heteroaliphatic moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, saturated or unsaturated, and each of the foregoing aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ and $-(\text{alkyl})\text{heteroaryl}$ moieties may be substituted or unsubstituted;

[0120] lviii) X^2 is $-\text{CH}(\text{Me})\text{Y}-$ where Y is $\left(\begin{array}{c} \text{O} \\ \parallel \\ \text{---} \text{N} \text{---} \text{CH} \text{---} \text{R}^{X2B} \\ | \\ \text{H} \end{array} \right)_2$ where each occurrence R^{X2B} is hydrogen, or an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, $-(\text{heteroalkyl})\text{aryl}$, or $-(\text{heteroalkyl})\text{heteroaryl}$ moiety; whereby each of the foregoing aliphatic and heteroaliphatic moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, saturated or unsaturated, and each of the foregoing aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ and $-(\text{alkyl})\text{heteroaryl}$ moieties may be substituted or unsubstituted;

[0121] lix) X^2 is absent;

[0122] lx) X^3 is $(\text{CHR}^{X3A})_k$, $-\text{CH}_2\text{NH}-$, $-\text{C}(=\text{O})\text{NH}-$, or $-\text{SO}_2-$, wherein each occurrence of R^{X3A} is independently hydrogen, an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, $-(\text{heteroalkyl})\text{aryl}$, or $-(\text{heteroalkyl})\text{heteroaryl}$ moiety, and k is an integer from 1 to 3 and each of the foregoing aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ and $-(\text{alkyl})\text{heteroaryl}$ moieties may be substituted or unsubstituted;


[0123] lxi) X^3 is absent;

[0124] lxii) X^3 is $-\text{C}(=\text{O})\text{NH}-$;

[0125] lxiii) X^3 is CHR^{X3A} , X^2 is $-\text{NR}^{X2A}-$ and R^{X2A} and R^{X3A} are each independently hydrogen or linear or branched substituted or unsubstituted alkyl;

[0126] lxiv) X^3 is CHR^{X3A} and X^2 is NH ; wherein R^{X3A} is hydrogen or linear or branched substituted or unsubstituted alkyl;


- [0127] lxxv) X^3 is CH_2 and X^2 is NH ;
- [0128] lxxvi) X^2 and X^3 are each absent;
- [0129] lxxvii) X^2 is CHR^{X2A} and X^3 is CH_2NH ; wherein R^{X2A} is hydrogen or linear or branched substituted or unsubstituted alkyl;
- [0130] lxxviii) X^2 is CHR^{X2A} and X^3 is $-\text{C}(=\text{O})\text{NH}-$; wherein R^{X2A} is hydrogen or linear or branched substituted or unsubstituted alkyl or $-\text{alkyl}(\text{aryl})$;
- [0131] lxxix) R^4 is substituted or unsubstituted, linear or branched, cyclic or acyclic alkyl, phenyl or $-(\text{CH}_2)\text{phenyl}$, wherein the phenyl group is optionally substituted with one or more occurrences of R^{4A} , wherein R^{4A} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ or $-(\text{alkyl})\text{heteroaryl}$, $-\text{OR}^{4B}$, $-\text{SR}^{4B}$, $-\text{N}(\text{R}^{4B})_2$, $-\text{SO}_2\text{N}(\text{R}^{4B})_2$, $-\text{C}(=\text{O})\text{N}(\text{R}^{4B})_2$, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OR}^{4B}$, $-\text{N}(\text{R}^{4B})\text{C}(=\text{O})\text{R}^{4C}$ or $-\text{N}(\text{R}^{4B})\text{SO}_2\text{R}^{4C}$; wherein each occurrence of R^{4B} and R^{4C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ or $-(\text{alkyl})\text{heteroaryl}$;
- [0132] lxxx) R^4 is substituted or unsubstituted, linear or branched, cyclic or acyclic alkyl, phenyl or $-(\text{CH}_2)\text{phenyl}$, wherein the phenyl group is optionally substituted with one or more occurrences of R^{4A} , wherein R^{4A} is hydrogen, hydroxyl, alkyl, alkoxy or halogen;
- [0133] lxxxi) R^4 is one of:



wherein each occurrence of R^{4A} is independently hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ or $-(\text{alkyl})\text{heteroaryl}$, $-\text{OR}^{4B}$, $-\text{SR}^{4B}$, $-\text{N}(\text{R}^{4B})_2$, $-\text{SO}_2\text{N}(\text{R}^{4B})_2$, $-\text{C}(=\text{O})\text{N}(\text{R}^{4B})_2$, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OR}^{4B}$, $-\text{N}(\text{R}^{4B})\text{C}(=\text{O})\text{R}^{4C}$, wherein each occurrence of R^{4B} and R^{4C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ or $-(\text{alkyl})\text{heteroaryl}$, wherein v and w are each independently integers from 0 to 3 and x is an integer from 1 to 6; whereby each of the foregoing alkyl and heteroalkyl moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, and each of the foregoing aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ and $-(\text{alkyl})\text{heteroaryl}$ moieties may be substituted or unsubstituted;

[0134] lxxii) R^4 is hydrogen;

[0135] lxxiii) R^4 is methyl, ethyl, propyl or one of:



wherein each occurrence of R^{4A} is independently hydrogen, lower alkyl or $C(=O)OR^{4B}$, wherein R^{4B} is hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl; each occurrence of X is independently a halogen; w is an integer from 0 to 3 and x is an integer from 1 to 6; whereby each of the foregoing alkyl and heteroalkyl moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, and each of the foregoing aryl, heteroaryl, -(alkyl)aryl and -(alkyl)heteroaryl moieties may be substituted or unsubstituted;

[0136] lxxiv) compounds of subset lxxiii) wherein w is 1;

[0137] lxxv) compounds of subset lxxiii) wherein w is 0;


[0138] lxxvi) compounds of subset lxxiii) wherein x is 1, 3 or 4;

[0139] lxxvii) compounds of subset lxxiii) wherein X is chlorine or fluorine;

[0140] lxxviii) compounds of subset lxxiii) wherein each occurrence of X is fluorine;


[0141] lxxix) compounds of subset lxxiii) wherein R^{4A} is methyl;

[0142] lxxx) R^4 is methyl, ethyl, propyl, isopropyl or one of:




wherein R^{4A} is hydrogen, hydroxyl, lower alkyl, lower alkoxy, halogen, $C(=O)OR^{4B}$, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl, wherein R^{4B} is hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl or -(heteroalkyl)heteroaryl;


[0143] lxxxi) R^4 is methyl, ethyl, propyl, isopropyl or one of:




[0144] lxxxii) R^4 is one of:



[0145] lxxxiii) R^4 is methyl, ethyl, propyl, isopropyl or one of:



[0146] lxxxiv) R^4 is methyl, ethyl, propyl or one of:



[0147] lxxxv) R' , R^3 and $R^{3'}$ are each hydrogen, X^1 is $-C(=O)-$, X^3 is $-C(=O)NH-$, R^1 is as described in subset xxiii, R^2 is as described in subset xxxiv, X^2 is as described in subset li, and R^4 is as described in subset lxxxiii;

[0148] lxxxvi) R' , R^3 and $R^{3'}$ are each hydrogen, X^1 is $-C(=O)-$, X^3 is $-C(=O)NH-$, R^1 is as described in subset xxiv, R^2 is as described in subset xxxv, X^2 is as described in subset lii, and R^4 is as described in subset lxxxiv; and/or

[0149] lxxxvii) R' , R^3 and $R^{3'}$ are each hydrogen, X^1 is $-C(=O)-$, X^3 is $-C(=O)NH-$, R^1 is as described in subset xxiv, R^2 is as described in subset xxxvi, X^2 is as described in subset lii, and R^4 is as described in subset lxxxiv.

[0150] It will be appreciated that for each of the classes and subclasses described above and herein, any one or more occurrences of aliphatic or heteroaliphatic may independently be substituted or unsubstituted, cyclic or acyclic, linear or branched, saturated or unsaturated and any one or more occurrences of aryl, heteroaryl, cycloaliphatic, cycloheteroaliphatic may be substituted or unsubstituted.

[0151] The reader will also appreciate that all possible combinations of the variables described in i)- through lxxxvii) above (*e.g.*, R', R¹, X¹, R², R³, R^{3'}, X², X³ and R⁴, among others) are considered part of the invention. Thus, the invention encompasses any and all compounds of formula I generated by taking any possible permutation of variables R', R¹, X¹, R², R³, R^{3'}, X², X³ and R⁴, and other variables/substituents (*e.g.*, R^{X1A}, R^{X2A}, Y, R^{X3A}, R^{4A}, etc.) as further defined for R', R¹, X¹, R², R³, R^{3'}, X², X³ and R⁴, described in i)- through lxxxvii) above.

[0152] For example, an exemplary combination of variables described in i)- through lxxxvii) above includes those compounds of Formula I wherein:

R' is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl;

R¹ is alkyl, heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl;

X¹ is -C(=O)-, -S(=O)-, -C(=NH)-, -C(=S)-, -NC(=O)-, -NC(=S)-, -N-C(=N-C≡N)-, -NS(O₂)-, -CHR^{X1A}-, -SO₂-, -COO-, -C(=O)C(R^{X1A})₂-, or -SC(=O)- wherein each occurrence of R^{X1A} is independently hydrogen, or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety;


R² is alkyl, heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl;

R³ is hydrogen, halogen, alkyl, heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl;

R^{3'} is hydrogen, halogen, or lower alkyl;

R⁴ is alkyl, heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl;

X² is absent, -NR^{X2A}-, -(CHR^{X2A})_j-, -NR^{X2A}Y-, or -(CHR^{X2A})_jY- wherein each occurrence of R^{X2A} is independently hydrogen or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, each occurrence of Y is independently




wherein, for each independent occurrence of t, R^{X2B} is hydrogen,

or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, and wherein each occurrence of j and t is independently an integer from 1 to 4; and


X^3 is absent, $-NHCO-$, $-NHSO_2-$, $-NHCONH-$, $-NHCOO-$, $-CH_2NH-$, $-C(=O)-$, $-S(=O)-$, $-C(=NH)-$, $-C(=S)-$, $-NC(=S)N-$, $-N-C(=N-C\equiv N)N-$, $-NS(O_2)N-$, $-SO_2-$, $-CONR^{X3A}-$, $-COO-$, $-(CHR^{X3A})_k-$, $-O-$, or $-NR^{X3A}-$, wherein each occurrence of R^{X3A} is independently hydrogen, an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, and k is an integer from 1 to 3.


[0153] Other exemplary combinations are illustrated by compounds of the following subgroups I and II:

[0154] I. Compounds having the structure (and pharmaceutically acceptable derivatives thereof):



wherein R^1 , R^2 , R^3 , $R^{3'}$, X^2 , X^3 and R^4 are as defined generally and in classes and subclasses herein.

In certain embodiments,  has one of the following structures:




wherein R^3 is hydrogen, halogen or an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(alkyl)aryl$, $-(alkyl)heteroaryl$, $-(heteroalkyl)aryl$, or $-(heteroalkyl)heteroaryl$ moiety;

R^4 is an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(alkyl)aryl$, $-(alkyl)heteroaryl$, $-(heteroalkyl)aryl$, or $-(heteroalkyl)heteroaryl$ moiety, or R^4 , taken together with $R^{X2A'}$ or a substituent present on Y or X^3 , may form a cycloaliphatic, cycloheteroaliphatic, aryl, or heteroaryl moiety;

R^{X2A} is hydrogen or an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(alkyl)aryl$, $-(alkyl)heteroaryl$, $-(heteroalkyl)aryl$, or $-(heteroalkyl)heteroaryl$ moiety;


$R^{X2A'}$ is hydrogen, or an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(alkyl)aryl$, $-(alkyl)heteroaryl$, $-(heteroalkyl)aryl$, or $-(heteroalkyl)heteroaryl$ moiety,

or $R^{X2A'}$ taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aryl or heteroaryl moiety;





Y is independently absent or is $\left(\begin{array}{c} \text{O} \\ \parallel \\ \text{---} \text{N} \text{---} \text{CH} \text{---} \text{R}^{X2B} \\ | \\ \text{H} \end{array} \right)_t$, wherein, for each independent occurrence of t , R^{X2B} is hydrogen, or an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, $-(\text{heteroalkyl})\text{aryl}$, or $-(\text{heteroalkyl})\text{heteroaryl}$ moiety, or one occurrence of R^{X2B} taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aryl or heteroaryl moiety; t is an integer from 1 to 4; and


X^3 is absent, $-\text{NHCO}-$, $-\text{NHSO}_2-$, $-\text{NHCONH}-$, $-\text{NHCOO}-$, $-\text{CH}_2\text{NH}-$, $-\text{C}(=\text{O})-$, $-\text{S}(=\text{O})-$, $-\text{C}(=\text{NH})-$, $-\text{C}(=\text{S})-$, $-\text{NC}(=\text{S})\text{N}-$, $-\text{N}-\text{C}(=\text{N}-\text{C}\equiv\text{N})\text{N}-$, $-\text{NS}(\text{O}_2)\text{N}-$, $-\text{SO}_2-$, $-\text{C}(=\text{O})\text{NR}^{X3A}-$, $-\text{C}(=\text{S})\text{NR}^{X3A}-$, $-\text{COO}-$, $(\text{CHR}^{X3A})_k$, $-\text{O}-$, $-\text{CH}_2\text{NR}^{X3A}-$, or $-\text{NR}^{X3A}-$, wherein each occurrence of R^{X3A} is independently hydrogen, an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, $-(\text{heteroalkyl})\text{aryl}$, or $-(\text{heteroalkyl})\text{heteroaryl}$ moiety, or R^{X3A} taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aryl or heteroaryl moiety, and k is an integer from 1 to 3.



[0155] In certain embodiments, $\begin{array}{c} \text{X}^2 \\ \diagup \quad \diagdown \\ \text{R}^3 \quad \text{R}^{3'} \quad \text{X}^3 \text{---} \text{R}^4 \end{array}$ has one of the following structures:




[0156] In certain embodiments, $\begin{array}{c} \text{X}^2 \\ \diagup \quad \diagdown \\ \text{R}^3 \quad \text{R}^{3'} \quad \text{X}^3 \text{---} \text{R}^4 \end{array}$ has one of the following structures:




wherein R^3 , R^4 , R^{X2A} , $R^{X2A'}$, R^{X2B} and t are as defined above and in classes and subclasses herein.

[0157] In certain embodiments, $\begin{array}{c} \text{X}^2 \\ \diagup \quad \diagdown \\ \text{R}^3 \quad \text{R}^{3'} \end{array} \text{X}^3\text{R}^4$ has one of the following structures:




wherein R^3 and R^4 are as defined above and in classes and subclasses herein.

[0158] II. Compounds having the structure (and pharmaceutically acceptable derivatives thereof):




wherein R^1 , R^2 , R^{X2A} and R^4 are as defined generally and in classes and subclasses herein.

[0159] In certain embodiments for compounds as described in subgroups I-II above, R^1 is a substituted or unsubstituted aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl or heterocyclic moiety. In certain exemplary embodiments, R^1 has one of the following structures:




[0160] In certain embodiments, for compounds as described in subgroups I-II above, R^2 has one of the following structures:



[0161] In certain embodiments, for compounds as described in subgroup I above, R^3 is halogen. In certain exemplary embodiments, R^3 is F. In certain other embodiments, R^3 is hydrogen.

[0162] In certain embodiments for compounds as described in subgroups I and II above, R^4 is substituted or unsubstituted, linear or branched, cyclic or acyclic alkyl, phenyl or $-(CH_2)phenyl$, wherein the phenyl group is optionally substituted with one or more occurrences of R^{4A} , wherein R^{4A} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, $-(alkyl)aryl$ or $-(alkyl)heteroaryl$, $-OR^{4B}$, $-SR^{4B}$, $-N(R^{4B})_2$, $-SO_2N(R^{4B})_2$, $-C(=O)N(R^{4B})_2$, halogen, $-CN$, $-NO_2$, $-C(=O)OR^{4B}$, $-N(R^{4B})C(=O)R^{4C}$, wherein each occurrence of R^{4B} and R^{4C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(alkyl)aryl$, $-(alkyl)heteroaryl$, $-(heteroalkyl)aryl$, $-(heteroalkyl)heteroaryl$. In certain other exemplary embodiments, R^4 is substituted or unsubstituted, linear or branched, cyclic or acyclic alkyl, phenyl or $-(CH_2)phenyl$, wherein the phenyl group is optionally substituted with one or more occurrences of R^{4A} , wherein R^{4A} is hydrogen, hydroxyl, alkyl, alkoxy or halogen. In certain exemplary embodiments, R^4 has one of the following structures:



[0163] In certain exemplary embodiments, for compounds as described in subgroups I and II above, R^{X2A} is a substituted or unsubstituted, linear or branched lower alkyl moiety. In certain other exemplary embodiments, R^{X2A} is methyl, ethyl, propyl, isopropyl or phenethyl.

[0164] It will also be appreciated that for each of the subgroups I-II described above, a variety of other subclasses are of special interest, including, but not limited to those classes described above i)-lxxxvii) and classes, subclasses and species of compounds described above and in the examples herein.

[0165] Some of the foregoing compounds can comprise one or more asymmetric centers, and thus can exist in various isomeric forms, *e.g.*, stereoisomers and/or diastereomers. Thus, inventive compounds and pharmaceutical compositions thereof may be in the form of an individual enantiomer, diastereomer or geometric isomer, or may be in the form of a mixture of stereoisomers. In certain embodiments, the compounds of the invention are enantiopure compounds. In certain other embodiments, mixtures of stereoisomers or diastereomers are provided.

[0166] Furthermore, certain compounds, as described herein may have one or more double bonds that can exist as either the Z or E isomer, unless otherwise indicated. The invention additionally encompasses the compounds as individual isomers substantially free of other isomers and alternatively, as mixtures of various isomers, *e.g.*, racemic mixtures of stereoisomers. In addition to the above-mentioned compounds *per se*, this invention also encompasses pharmaceutically acceptable derivatives of these compounds and compositions comprising one or more compounds of the invention and one or more pharmaceutically acceptable excipients or additives.

Compounds of the invention may be prepared by crystallization of compound of formula (I) under different conditions and may exist as one or a combination of polymorphs of compound of general formula (I) forming part of this invention. For example, different polymorphs may be identified and/or prepared using different solvents, or different mixtures of solvents for recrystallization; by performing

crystallizations at different temperatures; or by using various modes of cooling, ranging from very fast to very slow cooling during crystallizations. Polymorphs may also be obtained by heating or melting the compound followed by gradual or fast cooling. The presence of polymorphs may be determined by solid probe NMR spectroscopy, IR spectroscopy, differential scanning calorimetry, powder X-ray diffractogram and/or other techniques. Thus, the present invention encompasses inventive compounds, their derivatives, their tautomeric forms, their stereoisomers, their polymorphs, their pharmaceutically acceptable salts their pharmaceutically acceptable solvates and pharmaceutically acceptable compositions containing them.


[0167] **2) Synthetic Overview:**

[0168] The practitioner has a a well-established literature of peptide chemistry to draw upon, in combination with the information contained herein, for guidance on synthetic strategies, protecting groups, and other materials and methods useful for the synthesis of the compounds of this invention, including compounds containing the various R', R¹, R², R³, R^{3'} and R⁴ substituents and X¹, X² and X³ moieties.

[0169] The various patent documents and other references cited herein provide helpful background information on preparing compounds similar to the inventive compounds described herein or relevant intermediates, as well as information on formulation, uses, and administration of such compounds which may be of interest.

[0170] Moreover, the practitioner is directed to the specific guidance and examples provided in this document relating to various exemplary compounds and intermediates thereof.


[0171] As described above, the present invention provides novel compounds, specifically compounds having the following general structure:




and pharmaceutically acceptable derivatives thereof;

wherein R¹, R², R³, R^{3'}, R', X¹, X², X³ and R⁴ are as defined generally above and in classes and subclasses herein.


[0172] It will be appreciated that for compounds as generally described above, certain classes of compounds are of special interest. For example, one class of compounds of special interest includes those compounds wherein the compound has the stereochemistry as shown in Formula (I^A):



[0173] In yet another aspect of the invention, methods for producing intermediates useful for the preparation of compounds of formulae (I) and (I^A) are provided, embodiments of said methods being depicted generally in Schemes A and A₁:





Scheme A




Scheme A₁

[0174] Alternatively, substituent R³' may be introduced at a later stage in the synthesis, as depicted for example in Schemes A₂ and A₃:

Scheme A₂Scheme A₃

[0175] In certain embodiments, the inventive method comprises

i) providing a compound having the structure:



wherein R' , R^2 , R^3 , $R^{3'}$ and X^2 are as defined generally above and in classes and subclasses herein,


R^{P1} is a nitrogen protecting group;

A is absent, $-NHCO-$, $-NHSO_2-$, $-NHCONH-$, $-NHCOO-$, $-CH_2NH-$, $-C(=O)-$, $-S(=O)-$, $-C(=NH)-$, $-C(=S)-$, $-NC(=S)N-$, $-N-C(=N-C\equiv N)N-$, $-NS(O_2)N-$, $-SO_2-$, $-C(=O)NR^5-$, $-C(=S)NR^5-$, $-COO-$, $-(CHR^5)_r-$, $-O-$, $-CH_2NR^5-$ or $-NR^5-$, wherein each occurrence of R^5 is independently hydrogen, an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(alkyl)aryl$, $-(alkyl)heteroaryl$, $-(heteroalkyl)aryl$, or $-(heteroalkyl)heteroaryl$ moiety, and k is an integer from 1 to 3; and

B is a protecting group, or an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(alkyl)aryl$ or $-(alkyl)heteroaryl$ moiety, or is $-VR^C$, wherein V is $-O-$, $-NR^D-$, $-$


C(=O)-, -S(=O)- or -SO₂-, wherein each occurrence of R^C and R^D is independently hydrogen, a protecting group or an aliphatic, heteroaliphatic, aryl, heteroaryl, - (alkyl)aryl or -(alkyl)heteroaryl moiety;

ii) reacting the compound of step (i) under suitable conditions to generate a compound having the structure:




wherein U is -NHR^{P2} or -N₃, wherein R^{P2} is a nitrogen protecting group;

iii) reacting the compound of step (ii) with suitable reagents to generate a compound having the structure:




; and

iv) reacting the compound of step (iii) with suitable reagents to generate the free amine having the structure :




wherein R', R¹, R², R³, R^{3'}, R⁴, X¹, X² and X³ are as defined generally above and in classes and subclasses herein.

[0176] In certain embodiments, the present invention encompasses methods for the preparation of compounds having the general formula (I^A), and classes and subclasses herein.



[0177] In certain embodiments, the method comprises:

i) providing a compound having the structure:




wherein R' , R^2 , R^3 , $R^{3'}$ and X^2 are as defined above,

R^{P1} is a nitrogen protecting group;

A is absent, $-NHCO-$, $-NHSO_2-$, $-NHCONH-$, $-NHCOO-$, $-CH_2NH-$, $-C(=O)-$, $-S(=O)-$, $-C(=NH)-$, $-C(=S)-$, $-NC(=S)N-$, $-N-C(=N-C\equiv N)N-$, $-NS(O_2)N-$, $-SO_2-$, $-C(=O)NR^5-$, $-C(=S)NR^5-$, $-COO-$, $-(CHR^5)_r-$, $-O-$, $-CH_2NR^5-$ or $-NR^5-$, wherein each occurrence of R^5 is independently hydrogen, an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(alkyl)aryl$, $-(alkyl)heteroaryl$, $-(heteroalkyl)aryl$, or $-(heteroalkyl)heteroaryl$ moiety, and k is an integer from 1 to 3; and


B is a protecting group, or an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(alkyl)aryl$ or $-(alkyl)heteroaryl$ moiety, or is $-VR^C$, wherein V is $-O-$, $-NR^D-$, $-C(=O)-$, $-S(=O)-$ or $-SO_2-$, wherein each occurrence of R^C and R^D is independently hydrogen, a protecting group or an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(alkyl)aryl$ or $-(alkyl)heteroaryl$ moiety;

ii) reacting the compound of step (i) under suitable conditions to generate a compound having the structure:




wherein U is $-NHR^{P2}$ or $-N_3$, wherein R^{P2} is a nitrogen protecting group;

iii) reacting the compound of step (ii) with suitable reagents to generate a compound having the structure:



; and



iv) reacting the compound of step (iii) with suitable reagents to generate the free amine having the structure (I^A):



(I^A)


wherein R', R¹, R², R³, R^{3'}, R⁴, X¹, X² and X³ are as defined generally above and in classes and subclasses herein.

[0178] In yet another aspect of the invention, methods for producing intermediates useful for the preparation of compounds of formulas (I') and (I'^A) are provided, embodiments of said methods being depicted generally in Schemes A₄ and A₅:

Scheme A₄Scheme A₅

[0179] In certain embodiments, the inventive method comprises

i) providing a compound having the structure:



wherein R², R³, and X² are as defined above,


R^{P1} is a nitrogen protecting group;

A is absent, -NHCO-, -NHSO₂-, -NHCONH-, -NHCOO-, -CH₂NH-, -C(=O)-, -S(=O)-, -C(=NH)-, -C(=S)-, -NC(=S)N-, -N-C(=N-C≡N)N-, -NS(O₂)N-, -

SO₂-, -C(=O)NR⁵-, -C(=S)NR⁵-, -COO-, -(CHR⁵)_r-, -O-, -CH₂NR⁵- or -NR⁵-, wherein each occurrence of R⁵ is independently hydrogen, an aliphatic, heteroaliphatic, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety, and k is an integer from 1 to 3; and


B is a protecting group, or an aliphatic, heteroaliphatic, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl moiety, or is -VR^C, wherein V is -O-, -NR^D-, -C(=O)-, -S(=O)- or -SO₂-, wherein each occurrence of R^C and R^D is independently hydrogen, a protecting group or an aliphatic, heteroaliphatic, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl moiety;

ii) reacting the compound of step (i) under suitable conditions to generate a compound having the structure:




wherein U is -NHR^{P2} or -N₃, wherein R^{P2} is a nitrogen protecting group;

iii) reacting the compound of step (ii) with suitable reagents to generate a compound having the structure:




iv) reacting the compound of step (iii) with suitable reagents to generate the free amine having the structure :



wherein R¹, R², R³, R⁴, X¹, X² and X³ are as defined generally above and in classes and subclasses herein.


[0180] In certain embodiments, the present invention encompasses methods for the preparation of compounds having the general formula (I^A), and classes and subclasses herein.



(I^A)

[0181] In certain embodiments, the method comprises:

i) providing a compound having the structure:




wherein R^2 , R^3 , and X^2 are as defined above,

R^{P1} is a nitrogen protecting group;

A is absent, $-\text{NHCO}-$, $-\text{NHSO}_2-$, $-\text{NHCONH}-$, $-\text{NHCOO}-$, $-\text{CH}_2\text{NH}-$, $-\text{C}(=\text{O})-$, $-\text{S}(=\text{O})-$, $-\text{C}(=\text{NH})-$, $-\text{C}(=\text{S})-$, $-\text{NC}(=\text{S})\text{N}-$, $-\text{N}-\text{C}(=\text{N}-\text{C}\equiv\text{N})\text{N}-$, $-\text{NS}(\text{O}_2)\text{N}-$, $-\text{SO}_2-$, $-\text{C}(=\text{O})\text{NR}^5-$, $-\text{C}(=\text{S})\text{NR}^5-$, $-\text{COO}-$, $-(\text{CHR}^5)_r-$, $-\text{O}-$, $-\text{CH}_2\text{NR}^5-$ or $-\text{NR}^5-$, wherein each occurrence of R^5 is independently hydrogen, an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, $-(\text{heteroalkyl})\text{aryl}$, or $-(\text{heteroalkyl})\text{heteroaryl}$ moiety, and k is an integer from 1 to 3; and


B is a protecting group, or an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ or $-(\text{alkyl})\text{heteroaryl}$ moiety, or is $-\text{VR}^{\text{C}}$, wherein V is $-\text{O}-$, $-\text{NR}^{\text{D}}-$, $-\text{C}(=\text{O})-$, $-\text{S}(=\text{O})-$ or $-\text{SO}_2-$, wherein each occurrence of R^{C} and R^{D} is independently hydrogen, a protecting group or an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ or $-(\text{alkyl})\text{heteroaryl}$ moiety;

ii) reacting the compound of step (i) under suitable conditions to generate a compound having the structure:




wherein U is $-\text{NHR}^{\text{P2}}$ or $-\text{N}_3$, wherein R^{P2} is a nitrogen protecting group;

iii) reacting the compound of step (ii) with suitable reagents to generate a compound having the structure:




iv) reacting the compound of step (iii) with suitable reagents to generate the free amine having the structure (I^A):



wherein R¹, R², R³, R⁴, X¹, X² and X³ are as defined generally above and in classes and subclasses herein.

[0182] In certain exemplary embodiments, compounds (and methods of synthesis thereof) are provided wherein the compounds have the general formula (I^{''A}) as shown in Scheme B below:



Scheme B


wherein each of R¹, X¹, R^{P1}, R², X³, and R⁴ are as defined above and R^{P3} is an oxygen protecting group.

[0183] It will be appreciated that in certain embodiments, as depicted in Scheme B, step i) involves providing a oxo-tetrahydro-furan-2-yl (as described in

the Examples herein). Subsequent ring opening, protection, oxidation and stereoselective reduction yields the desired compound of provided in step i). Subsequent reduction to an azide in step ii) and reaction with a suitable reagent (to yield $-X^1-R^1$) is also effected. Finally, reaction with a suitable reagent to generate $-X^3-R^4$ and deprotection yields the desired compound. It will be appreciated that a variety of methods can be utilized to effect these transformations, including those detailed in the specification herein, and additional general guidance as also described herein.


[0184] Additionally, it will be appreciated that compounds having the general structures depicted above, wherein X^3 is SO_2 can be effected using the methodology generally described herein and also methodology described in 5,585,397, the entire contents of which are hereby incorporated by reference.

[0185] Numerous suitable prodrug moieties, and information concerning their selection, synthesis and use are well known in the art. Examples of prodrug moieties of interest include, among others, prodrug moieties that can be attached to primary or secondary amine-containing functionalities. For instance, prodrug moieties of interest include those that can be attached to group $-NH_2$. Examples of such prodrug moieties include the following:




R^1 = all natural,
unnatural amino acids

For the synthesis of the prodrug groups, see Borchardt, R. T. et. al., *J. Org. Chem.* 1997, 43, 3641-3652.



R^1 = C1-C4 alkyl, cycloalkyl, oxyalkyl,
aminoalkyl, etc.
 R^2 = all natural, unnatural amino acids

For the synthesis of the prodrug groups, see Zhou, X-X. et. al., PCT WO 99/51613.



R^1, R^2 = all natural, unnatural amino acids

For the synthesis of the prodrug groups, see Ezra, A. et. al., *J. Med. Chem.* 2000, 43, 3641-3652.

[0186] The present invention encompasses any prodrug form of the compounds described herein. Although certain other exemplary prodrug moieties generated from the inventive compounds amino group are detailed herein, it will be appreciated that the present invention is not intended to be limited to these prodrug moieties; rather, a variety of additional prodrug moieties can be readily identified by a person skilled in the relevant art.

[0187] **3) Pharmaceutical Compositions**

[0188] As discussed above, certain of the compounds as described herein exhibit activity generally as inhibitors of aspartyl proteases and more specifically as inhibitors of β -secretase enzyme activity, and have the ability to halt or reduce the production of A β from APP and reduce or eliminate the formation of β -amyloid deposits in the brain. Thus, the compounds are useful for treating humans or animals suffering from a condition characterized by a pathological form of β -amyloid peptide, such as β -amyloid plaques, and for helping to prevent or delay the onset of such a condition. Thus, in certain embodiments, compounds of the invention are useful for treating Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's disease, for treating patients with MCI (mild cognitive impairment) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, *i.e.* single and recurrent lobar hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, and diffuse Lewy body type Alzheimer's disease. The compounds and compositions of the invention are particularly useful for treating or preventing Alzheimer's disease. When treating or preventing these diseases, the compounds of the invention can either be used individually or in combination.

[0189] Accordingly, in another aspect of the present invention, pharmaceutical compositions are provided, which comprise any one of the compounds described herein (or a prodrug, pharmaceutically acceptable salt or other

pharmaceutically acceptable derivative thereof), and optionally comprise a pharmaceutically acceptable carrier. In certain embodiments, these compositions optionally further comprise one or more additional therapeutic agents. Alternatively, a compound of this invention may be administered to a patient in need thereof in combination with the administration of one or more other therapeutic agents. For example, additional therapeutic agents for conjoint administration or inclusion in a pharmaceutical composition with a compound of this invention may be an approved agent for the treatment of Alzheimer's Disease, or it may be any one of a number of agents undergoing approval in the Food and Drug Administration that ultimately obtain approval for the treatment any disorder suffering from a condition characterized by a pathological form of β -amyloid peptide. It will also be appreciated that certain of the compounds of present invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable derivative thereof. According to the present invention, a pharmaceutically acceptable derivative includes, but is not limited to, pharmaceutically acceptable salts, esters, salts of such esters, or a pro-drug or other adduct or derivative of a compound of this invention which upon administration to a patient in need is capable of providing, directly or indirectly, a compound as otherwise described herein, or a metabolite or residue thereof.

[0190] As used herein, the term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts of amines, carboxylic acids, and other types of compounds, are well known in the art. For example, S.M. Berge, *et al.* describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 66: 1-19 (1977), incorporated herein by reference. The salts can be prepared *in situ* during the final isolation and purification of the compounds of the invention, or separately by reacting a free base or free acid function with a suitable reagent, as described generally below. For example, a free base function can be reacted with a suitable acid. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may,

include metal salts such as alkali metal salts, *e.g.* sodium or potassium salts; and alkaline earth metal salts, *e.g.* calcium or magnesium salts. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, harnisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

[0191] Additionally, as used herein, the term “pharmaceutically acceptable ester” refers to esters that hydrolyze *in vivo* and include those that break down readily in the human body to leave the parent compound or a salt thereof. Suitable ester groups include, for example, those derived from pharmaceutically acceptable aliphatic carboxylic acids, particularly alkanoic, alkenoic, cycloalkanoic and alkanedioic acids, in which each alkyl or alkenyl moiety advantageously has not more than 6 carbon atoms. Examples of particular esters include formates, acetates, propionates, butyrates, acrylates and ethylsuccinates.

[0192] Furthermore, the term “pharmaceutically acceptable prodrugs” as used herein refers to those prodrugs of the compounds of the present invention

which are, within the scope of sound medical judgment, suitable for use in contact with the issues of humans and lower animals with undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term "prodrug" refers to compounds that are rapidly transformed *in vivo* to yield the parent compound of the above formula, for example by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference.

[0193] As described above, the pharmaceutical compositions of the present invention additionally comprise a pharmaceutically acceptable carrier, which, as used herein, includes any and all solvents, diluents, or other liquid vehicle, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington's Pharmaceutical Sciences, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1980) discloses various carriers used in formulating pharmaceutical compositions and known techniques for the preparation thereof. Except insofar as any conventional carrier medium is incompatible with the compounds of the invention, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutical composition, its use is contemplated to be within the scope of this invention. Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatine; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil, sesame oil; olive oil; corn oil and soybean oil; glycols; such as propylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogenfree water;

isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

[0194] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[0195] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[0196] The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[0197] In order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension or crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle. Injectable depot may form are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include (poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

[0198] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

[0199] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium

stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[0200] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[0201] The active compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose and starch. Such dosage forms may also comprise, as in normal practice, additional substances other than inert diluents, *e.g.*, tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[0202] Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed

under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, eardrops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. These dosage may form are made by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[0203] It will also be appreciated that the compounds and pharmaceutical compositions of the present invention can be formulated and employed in combination therapies, that is, the compounds and pharmaceutical compositions can be formulated with or administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. For example, compounds of the invention can be used in combination, with each other, or with other therapeutic agents or approaches used to treat or prevent the conditions described above and herein. Exemplary agents include, but are not limited to: acetylcholine esterase inhibitors such as tacrine (tetrahydroaminoacridine, marketed as COGNEX®), donepezil hydrochloride (marketed as Aricept® and rivastigmine (marketed as Exelon®); gamma-secretase inhibitors; anti-inflammatory agents such as cyclooxygenase II inhibitors; anti-oxidants such as Vitamin E and ginkgolides; immunological approaches, such as, for example, immunization with A β peptide or administration of anti-A β peptide antibodies; statins; and direct or indirect neurotropic agents such as Cerebrolysin®, AIT-082 (Emilieu, 2000, *Arch. Neurol.* 57:454), and other neurotropic agents of the future. The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed may achieve a desired effect for the same disorder (for example, an inventive compound may be administered concurrently with another Alzheimer's agent), or they may achieve different effects (*e.g.*, control of any adverse effects). In

certain embodiments, the pharmaceutical compositions of the present invention further comprises one or more additional therapeutically active ingredients (*e.g.*, palliative). For purposes of the invention, the term “*Palliative*” refers to treatment that is focused on the relief of symptoms of a disease and/or side effects of a therapeutic regimen, but is not curative. For example, palliative treatment encompasses painkillers, antinausea medications and anti-sickness drugs.

[0204] 4) Research Uses, Pharmaceutical Uses and Methods of Treatment

[0205] *Research Uses*

[0206] According to the present invention, the inventive compounds may be assayed in any of the available assays known in the art for identifying compounds having protease inhibitory activity. For example, the assay may be cellular or non-cellular, *in vivo* or *in vitro*, high- or low-throughput format, etc.

[0207] In certain exemplary embodiments, compounds of this invention were assayed for their ability to inhibit aspartyl proteases, more specifically BACE.

[0208] Thus, in one aspect, compounds of this invention which are of particular interest include those which:

- are inhibitors of aspartyl proteases;
- exhibit the ability to inhibit BACE (β -secretase enzyme activity);
- exhibit the ability to inhibit A β peptide production;
- exhibit the ability to halt or reduce the production of A β from APP and reduces or eliminates the formation of β -amyloid deposits in the brain;
- are useful for treating mammals (*e.g.*, humans) or animals suffering from a condition characterized by a pathological form of β -amyloid peptide, such as β -amyloid plaques, and for helping to prevent or delay the onset of such a condition;
- exhibit a favorable therapeutic profile (*e.g.*, safety, efficacy, and stability).

[0209] In certain embodiments, compounds of the invention are aspartyl protease inhibitors. In certain exemplary embodiments, inventive compounds are selective BACE inhibitors. In certain exemplary embodiments, inventive compounds have $\text{BACE}K_i^{\text{app}} \leq 10 \mu\text{M}$. In certain other embodiments, inventive compounds have $\text{BACE}K_i^{\text{app}} \leq 7.5 \mu\text{M}$. In certain other embodiments, inventive compounds have

$BACE_{K_i^{app}} \leq 5 \mu M$. In certain other embodiments, inventive compounds have $BACE_{K_i^{app}} \leq 2.5 \mu M$. In certain other embodiments, inventive compounds have $BACE_{K_i^{app}} \leq 1 \mu M$. In certain other embodiments, inventive compounds have $BACE_{K_i^{app}} \leq 750 \text{ nM}$. In certain other embodiments, inventive compounds have $BACE_{K_i^{app}} \leq 500 \text{ nM}$. In certain other embodiments, inventive compounds have $BACE_{K_i^{app}} \leq 250 \text{ nM}$. In certain other embodiments, inventive compounds have $BACE_{K_i^{app}} \leq 100 \text{ nM}$. In certain other embodiments, inventive compounds have $BACE_{K_i^{app}} \leq 80 \text{ nM}$. In certain other embodiments, inventive compounds have $BACE_{K_i^{app}} \leq 60 \text{ nM}$. In certain other embodiments, inventive compounds have $BACE_{K_i^{app}} \leq 50 \text{ nM}$. In certain other embodiments, inventive compounds have $BACE_{K_i^{app}} \leq 30 \text{ nM}$. In certain other embodiments, inventive compounds have $BACE_{K_i^{app}} \leq 20 \text{ nM}$. In certain other embodiments, inventive compounds have $BACE_{K_i^{app}} \leq 10 \text{ nM}$.

[0210] In certain embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 2 fold greater than $BACE_{K_i^{app}}$. In certain other embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 3 fold greater than $BACE_{K_i^{app}}$. In certain other embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 4 fold greater than $BACE_{K_i^{app}}$. In certain other embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 5 fold greater than $BACE_{K_i^{app}}$. In certain embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 7.5 fold greater than $BACE_{K_i^{app}}$. In certain embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 10 fold greater than $BACE_{K_i^{app}}$. In certain embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 25 fold greater than $BACE_{K_i^{app}}$. In certain embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 50 fold greater than $BACE_{K_i^{app}}$. In certain embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 75 fold greater than $BACE_{K_i^{app}}$. In certain embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 100 fold greater than $BACE_{K_i^{app}}$. In certain embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 150 fold greater than $BACE_{K_i^{app}}$. In certain embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 200 fold greater than $BACE_{K_i^{app}}$. In certain embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 250 fold greater than $BACE_{K_i^{app}}$. In certain embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 300 fold greater than $BACE_{K_i^{app}}$. In certain embodiments, $CatD_{K_i^{app}}$ for compounds of the invention is ≥ 500 fold greater than $BACE_{K_i^{app}}$. In certain

embodiments, $\text{CatD}K_i^{\text{app}}$ for compounds of the invention is ≥ 1000 fold greater than $\text{BACE}K_i^{\text{app}}$.

[0211] In certain exemplary embodiments, inventive compounds have CellIC_{50} values $\leq 10 \mu\text{M}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 7.5 \mu\text{M}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 5 \mu\text{M}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 2.5 \mu\text{M}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 1 \mu\text{M}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 750 \text{ nM}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 500 \text{ nM}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 250 \text{ nM}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 100 \text{ nM}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 80 \text{ nM}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 60 \text{ nM}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 50 \text{ nM}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 30 \text{ nM}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 20 \text{ nM}$. In certain other embodiments, inventive compounds have CellIC_{50} values $\leq 10 \text{ nM}$.

[0212] *Pharmaceutical Uses and Methods of Treatment*

[0213] As discussed above, without wishing to be bound by any particular theory, certain of the compounds as described herein exhibit activity generally as inhibitors of aspartyl proteases and more specifically as inhibitors of β -secretase enzyme activity. In certain embodiments compounds exhibit the ability to halt or reduce the production of A β from APP and reduce or eliminate the formation of β -amyloid deposits in the brain and thus the compounds are useful for treating humans or animals suffering from a condition characterized by a pathological form of β -amyloid peptide, such as β -amyloid plaques, and for helping to prevent or delay the onset of such a condition (*e.g.*, Alzheimer's Disease). Thus, in certain embodiments, compounds of the invention are useful for treating Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's disease, for treating patients with MCI (mild cognitive impairment) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating

Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, *i.e.* single and recurrent lobal hemorrhages, for treating other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, and diffuse Lewy body type Alzheimer's disease. The compounds and compositions of the invention are particularly useful for treating or preventing Alzheimer's disease. When treating or preventing these diseases, the compounds of the invention can either be used individually or in combination.

[0214] In certain embodiments, the method involves the administration of a therapeutically effective amount of the compound or a pharmaceutically acceptable derivative thereof to a subject (including, but not limited to a human or animal) in need of it.

[0215] As described in more detail herein, in certain embodiments, compounds of the invention are useful as inhibitors of aspartyl proteases. Generally, compounds of the invention exhibit the ability to inhibit aspartyl protease enzyme activity and thus compounds of the invention generally are useful for the treatment of disorders mediated by aspartyl protease enzyme activity. More specifically, compounds of the invention exhibit activity as inhibitors of β -secretase enzyme activity and A β peptide production. Thus, in certain embodiments, the present invention provides compounds useful for the treatment of disorders mediated by a pathological form of β -amyloid peptide, such as β amyloid plaques, and for helping to prevent or delay the onset of such a condition.

[0216] Thus, in certain embodiments, compounds of the invention are useful for treating Alzheimer's disease, for helping prevent or delay the onset of Alzheimer's disease, for treating patients with MCI (mild cognitive impairment) and preventing or delaying the onset of Alzheimer's disease in those who would progress from MCI to AD, for treating Down's syndrome, for treating humans who have Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, for treating cerebral amyloid angiopathy and preventing its potential consequences, *i.e.* single and recurrent lobal hemorrhages, for treating other degenerative dementias,

including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, and diffuse Lewy body type Alzheimer's disease. The compounds and compositions of the invention are particularly useful for treating or preventing Alzheimer's disease. When treating or preventing these diseases, the compounds of the invention can either be used individually or in combination.

[0217] Thus, as described above, in another aspect of the invention, a method for the treatment of disorders useful for the treatment (or prevention) of disorders mediated by a pathological form of β -amyloid peptide, such as β amyloid plaques, is provided comprising administering a therapeutically effective amount of a compound of Formula (I) or (I^A), or any classes and subclasses of these compounds as described herein, to a subject in need thereof. It will be appreciated that the compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for the treatment of disorders by a pathological form of β -amyloid peptide, such as β amyloid plaques. Thus, the expression "effective amount" as used herein, refers to a sufficient amount of agent to inhibit the production of A β peptide, and to exhibit a therapeutic effect. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular therapeutic agent, its mode of administration, and the like. The compounds of the invention are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression "dosage unit form" as used herein refers to a physically discrete unit of therapeutic agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and

rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts (see, for example, Goodman and Gilman's, "The Pharmacological Basis of Therapeutics", Tenth Edition, A. Gilman, J. Hardman and L. Limbird, eds., McGraw-Hill Press, 155-173, 2001, which is incorporated herein by reference in its entirety).

[0218] Another aspect of the invention relates to a method for inhibiting β -secretase activity in a biological sample or a patient, which method comprises administering to the patient, or contacting said biological sample with a compound of formula I or a composition comprising said compound.

[0219] Furthermore, after formulation with an appropriate pharmaceutically acceptable carrier in a desired dosage, the pharmaceutical compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, as an oral or nasal spray, or the like, depending on the severity of the infection being treated. In certain embodiments, the compounds of the invention may be administered at dosage levels of about 0.001 mg/kg to about 50 mg/kg, from about 0.01 mg/kg to about 25 mg/kg, or from about 0.1 mg/kg to about 10 mg/kg of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect. It will also be appreciated that dosages smaller than 0.001 mg/kg or greater than 50 mg/kg (for example 50-100 mg/kg) can be administered to a subject. In certain embodiments, compounds are administered orally or parenterally.

TREATMENT KIT

[0220] In other embodiments, the present invention relates to a kit for conveniently and effectively carrying out the methods in accordance with the present invention. In general, the pharmaceutical pack or kit comprises one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Such kits are especially suited for the delivery of solid oral forms such as tablets or capsules. Such a kit preferably includes a number of unit dosages, and may also include a card having the dosages oriented in the order

of their intended use. If desired, a memory aid can be provided, for example in the form of numbers, letters, or other markings or with a calendar insert, designating the days in the treatment schedule in which the dosages can be administered. Alternatively, placebo dosages, or calcium dietary supplements, either in a form similar to or distinct from the dosages of the pharmaceutical compositions, can be included to provide a kit in which a dosage is taken every day. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceutical products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

EQUIVALENTS

[0221] The representative examples that follow are intended to help illustrate the invention, and are not intended to, nor should they be construed to, limit the scope of the invention. Indeed, various modifications of the invention and many further embodiments thereof, in addition to those shown and described herein, will become apparent to those skilled in the art from the full contents of this document, including the examples which follow and the references to the scientific and patent literature cited herein. It should further be appreciated that the contents of those cited references are incorporated herein by reference to help illustrate the state of the art.

[0222] The following examples contain important additional information, exemplification and guidance that can be adapted to the practice of this invention in its various embodiments and the equivalents thereof.

EXEMPLIFICATION

[0223] The compounds of this invention and their preparation can be understood further by the examples that illustrate some of the processes by which these compounds are prepared or used. It will be appreciated, however, that these examples do not limit the invention. Variations of the invention, now known or further developed, are considered to fall within the scope of the present invention as described herein and as hereinafter claimed.

[0224] EXAMPLE 1

[0225] This example describes the enzyme assay for determining the apparent K_i of the compounds of the present invention.

[0226] The BACE enzyme used for inhibition analyses, BACE-HT, was produced from baculovirus-infected insect cells, and corresponded to the pro form of the soluble N-terminal protease domain (residues 22-454, starting from the N-terminal methionine), followed by a short linker and C-terminal 6X His tag.

[0227] Compounds were tested for their ability to inhibit BACE hydrolysis of the internally quenched fluorescent substrate, FS-1 (Ermolieff *et al.*, Biochemistry 39:12450-12456 (2000)):

FS1: $\text{NH}_2\text{-Arg-Glu(EDANS)-Glu-Val-Asn-Leu-}\downarrow\text{-Asp-Ala-Glu-Phe-Lys(DABCYL)-Arg-COOH}$

FS-2: $\text{MOCac-Ser-Glu-Val-Asn-Leu-}\downarrow\text{-Asp-Ala-Glu-Phe-Lys(DNP)-Arg-Arg-COOH}$

where EDANS is 5-((2-aminoethyl)amino)naphthalene-1-sulfonic acid; DABCYL is 4-((4-dimethylamino)phenyl)azo)benzoic acid; MOCac is 7-methoxycoumarin-4-yl) acetic acid; and DNP is 2,4-dinitrophenylacetic acid.

[0228] FS-1 and FS-2 correspond to the Swedish APP β -site sequence, flanked by either an EDANS or MOCac fluorophore on the N-terminus and a DABCYL or DNP quenching group near the C-terminus. In the intact state, the DABCYL or DNP group quenches the EDANS or MOCac (respectively) fluorescence by virtue of their close proximity. In the intact state, the DABCYL group quenches the EDANS fluorescence by virtue of their close proximity. Upon cleavage at the site indicated by the arrow by BACE, the quenching is relieved and fluorescence is observed using $\lambda_{\text{excitation}} = 350 \text{ nm}$ and $\lambda_{\text{emission}} = 490 \text{ nm}$ or $\lambda_{\text{excitation}} = 328 \text{ nm}$ and $\lambda_{\text{emission}} = 440 \text{ nm}$ (MOCac). For determination of apparent inhibition constant (K_i^{app}) for various inhibitors, the initial rates of FS-1 hydrolysis in the presence of various concentrations of inhibitors were measured and fit to the Morrison equation (Williams and Morrison, Methods Enzymol. 63: 437-467 (1979)):

$$v = v_0 \cdot \frac{[E]_0 - [I]_0 - K_i^{\text{app}} + \sqrt{([E]_0 - [I]_0 - K_i^{\text{app}})^2 + 4[E]_0 K_i^{\text{app}}}}{2[E]_0}$$

where v is the initial rate measured in the presence of $[I]_0$, the inhibitor concentration, using an enzyme concentration $[E]_0$. v_0 is the initial rate measured in the absence of inhibitor.

[0229] Inhibitors were resuspended in DMSO and serially diluted in DMSO at 20x final assay concentration. Compound dilutions (5 μ L) were transferred to Black non-treated 96-well microtiter plates, and resuspended in 85 μ L 35.3 μ M FS1 substrate in 100 mM sodium acetate buffer, pH 4.5, containing 5.9% DMSO (*i.e.* 10% final DMSO in assay). Following brief equilibration to room temperature, the reactions were initiated by the addition of 10 μ L BACE-HT (100nM final concentration) and brief mixing. The increase in EDANS fluorescence over time was monitored on a Gemini XS fluorometric plate reader using Softmax pro software. Initial rates were fit to the Morrison equation and K_i^{app} 's determined using Graphpad Prism software. A representative range of K_i^{app} 's for the compounds is between 1-1000 nM. The range of K_i^{app} 's for preferred compounds is 0.1-500 nM.

[0230] EXAMPLE 2

[0231] This example describes an illustrative set of cell-based assays for testing the ability of the compounds of the present invention to inhibit the secretion of the A β peptide from cells expressing high levels of the human amyloid precursor protein ("APP").

[0232] A representative set of such cells include A-204 rhabdomyosarcoma cells, human embryonic kidney 293 cells transiently transfected with APP, Chinese hamster ovary cells stably transfected with APP (CHO2B7), and H4 neuroglioma cells stably transfected with APP (H4 β 695wt). The A-204 cells and HEK 293 cells were obtained from ATCC. For HEK 293 transient transfections, a pcDNA3 plasmid containing the human APP gene was obtained from Invitrogen, and cells were transfected with FuGENE 6 transfection reagent (Roche) following manufacturer's protocols. The CHO2B7 and H4 β 695wt lines were obtained from Mayo Clinic (*J. Neuroscience Methods* 108(2): 171-9 (2001)).

[0233] The following media were used for the cell lines: DMEM + glucose, glutamine, sodium pyruvate, pyridoxine-HCl, antibiotics, and 10% FBS (A-204 cells); DMEM + glucose, glutamine, sodium pyruvate, pyridoxine-HCl, non-essential amino acids, antibiotics, and 10% FBS (HEK 293 cells); Ham's F-12 + antibiotics, 10% FBS, and 400 µg/ml Zeocin (CHO2B7 cells); and Opti-MEM (Invitrogen)+ antibiotics, 10% FBS, and 500 µg/ml geneticin(H4 β695wt cells).

[0234] In the assay, cells were pre-treated with compounds for 1-2 hours to "wash out" any Aβ in the secretory pathway prior to incubation with compounds for the conditioning period required to generate detectable concentrations of Aβ. The production of secreted Aβ in cultured cells was monitored by sandwich ELISA, using the anti-Aβ monoclonal antibodies 6E10 and 4G8 (Signet Laboratories). The antibody 6E10 (specific to amino acids 1-17 of Aβ) was immobilized to the plate and used for capture, and a biotinylated version of the antibody 4G8 (specific to amino acids 17-24 of Aβ) was used for detection, via subsequent treatment with Neutravidin-horseradish peroxidase conjugate and development with luminescent substrate.

[0235] Cells were plated at 2.5×10^5 cells/well (HEK293 cells in 24 well poly-D-lysine coated plate), 1.3×10^5 cells/well (A-204 cells) (48 well plates) or 5×10^4 cells/well (CHO2B7 and H4APP cells) (96 well plates) in appropriate media (500 µL per well for 24 well plates, 300 µL per well for 48 well plates, 150 µL per well for 96 well plates) and grown at 37° C and 5% CO₂ for 16-20 h. Media was aspirated and replaced with appropriate media plus 0.1% DMSO and various concentrations of inhibitors, prepared by resuspension of the appropriate volume of a 1000X solution of inhibitor in DMSO with media on a separate plate. Cells and inhibitors were incubated for 1-2 hours, followed by aspiration of the media and replacement with fresh media solutions having the same concentrations of inhibitors. After a 4-24 hour incubation period (depending on cell type), conditioned media were collected and 100 µL of each sample applied to wells of Nunc Maxisorp white 96-well plates that had been previously coated with 10 µg/mL 6E10 antibody in 50 mM Na₂CO₃, (pH 9.0) overnight at 4° C, blocked with Superblock in PBS (Pierce Chemicals; PBS=10 mM sodium phosphate + 150 mM NaCl, pH 7.4) for 1 hour at room temperature, and washed 3 times in wash buffer (PBS plus 0.05% Tween 20).

Binding was allowed to take place overnight at 4 °C, and plates were then washed 8 times and treated with 100 µL/well of 10 nM biotinylated 4G8 antibody in binding buffer (superblock in PBS plus 0.05% Tween 20) for 1 hour at room temperature. Plates were washed 8 times, treated with 100 µL/well of 1.5 µg/ Neutravidin-horseradish peroxidase conjugate (Pierce Chemicals) for 40 minutes at room temperature, washed an additional 8 times, and developed with 100 µL/well Supersignal luminescent substrate (Pierce Chemicals) for 2 minutes. Luminescence was measured on a Gemini XS fluorescence/luminescence plate reader, and translated into Aβ concentrations using an internal standard curve of known Aβ concentrations (Aβ 1-40 obtained from Oncogene). In general, IC₅₀'s of the compounds of the present invention range from about 5X to about 15X the observed K_i^{app}'s against the BACE enzyme as determined using the method of Example 1.

[0236] EXAMPLE 3

[0237] This example describes permeability experiments using MDCK cells.

[0238] MDCK cells, derived from canine kidney, were obtained from American Type Culture Collection (ATCC, Rockville, MD). Cells were cultured in DMEM medium supplemented with 10% fetal bovine serum, 1 mM sodium pyruvate and 0.01 mg/ml gentamicin. Cells were maintained in a humidified atmosphere with 5% CO₂ at 37°C. For transport studies, cells were plated at a density of 50 x 10³ cells/cm² on 0.4 µm pore size Transwell™ polyester membranes (Corning, Corning, NY). Culture medium was replaced every two days until a tight cell monolayer was formed as measured by preliminary R123 permeability measurements. Compounds, 10 µM, were placed in either the apical or basolateral compartment and the amount of drug in the opposite chamber was measured using LC/MS/MS or fluorometry. The permeability coefficient was calculated using the following equation:

$$P_{app} = (1/AC_0) dQ/dt$$

where A= surface area of the cell monolayers, C₀ is the initial concentration and dQ/dt represents the amount of drug flux in a specified time.

[0239] EXAMPLE 4


[0240] This example describes pharmacokinetic experiments for determining peak plasma and peak brain concentrations using Adult Swiss Weber mice. The peak brain concentration determination also served as a measure for assessing whether the compounds were able to cross the blood brain barrier.

[0241] Adult Swiss Webster mice were administered compounds via tail vein injection.

Plasma and brain tissue were collected at 0, 0.5, 1, 2, 3, 4, and 8 hrs. Blood was collected via cardiac puncture. Subsequently, animals were perfused intra-cardially with ice-cold saline; the brain was homogenized in 10 volumes of ice-cold Tris-buffered saline (20 mM Tris, 137 mM NaCl, pH 7.6) and was then frozen at -20°C . Following extraction, compound concentrations in plasma and brain homogenates were measured by LC/MS/MS. Peak plasma and brain concentrations (C_{max}) and time to achieve these concentrations (T_{max}) were measured directly from concentration versus time profiles. Descriptive pharmacokinetic parameters were calculated using the WinNonlin software package (Pharsight Inc., Mountain View CA). Figure 1A and 1B are the plasma and brain concentration curves respectively for one of the compounds of the present invention (where each data point is an average concentration from three mice). This compound has a molecular weight of 626, has a K_i^{app} of about 35 nM with a 125 fold selectivity over Cat D, and has an IC_{50} in cells of about 220 nM.

[0242] EXAMPLE 5

[0243] This example compares the biological properties of a hydroxy-containing compound with an amino-containing compound. The two compounds are of the structure



where R is either OH or NH₂. In a surprising and unexpected finding, the amino-containing compound possesses several superior biological properties over its hydroxy-containing counterpart.

[0244] As it can be seen in the following Table, the amino compound is more potent in cells, is more selective for the BACE enzyme, and show superior ADME properties than its hydroxy counterpart.

Compound	BACE K _i	IC ₅₀ (H4 cells)	Cat D K _i	% recovery liver microsomes
R = OH	150 nM	2000 nM	20 nM	0% at 60 minutes
R = NH ₂	180 nM	740 nM	610 nM	43% at 60 minutes

[0245] Although both compounds show similar activity against the BACE enzyme (180 nM versus 150 nM), the amino compound is significantly more potent in cells (740 nM versus 2000 nM). Moreover, the amino compound displays over a three fold preference for BACE over Cat D. In contrast, the hydroxy counterpart displays a 7.5 fold preference for Cat D over BACE. In addition, although both the amino and hydroxy compounds were stable in human plasma (100% recovery at 1 hour), the amino compound is much more stable in liver microsomes than its hydroxy counterpart.

[0246] These trends appear to be a general phenomenon and have been observed in several scaffolds in addition to one exemplified above.

[0247] The K_i for BACE and the IC₅₀ determinations were performed as described in Examples 1 and 2 respectively.

[0248] The K_i determination for Cat D was performed using the assay conditions derived from Haque *et al.*, *J. Med. Chem.* **42**: 1428-1440 (1999) and the following substrate:

DABCYL-GLu-Arg-Nle-Phe-Leu-Ser-Phe-Pro-EDANS where Nle is norleucine and DABCYL and EDANS are as previously defined.

[0249] Plasma stability was determined by incubating the compounds in human plasma at a concentration of 1 μM at 37 °C for 0, 30 and 60 minutes. Reactions were stopped by addition of acetonitrile. Protein was precipitated by centrifugation (3000 rpm x 10 min) then supernatants were analyzed for remaining parent compound by LC/MS/MS.

[0250] Metabolic stability was determined by incubating the compounds with human liver microsomes (100 µg/ml) in 100 mM Tris buffer pH 7.4 for 5 minutes at 37 °C. Metabolic reactions were started by addition of NADPH to give a final concentration of 1 µM. Reactions were stopped after 30 and 60 minutes by addition of acetonitrile. Protein was precipitated by centrifugation (3000 rpm x 10 min) then supernatants were analyzed for remaining parent compound by LC/MS/MS.

[0251] *General Description of Synthetic Strategy:*

[0252] According to the present invention, any available techniques can be used to make or prepare the inventive compounds or compositions including them. For example, a variety of solution phase synthetic methods such as those discussed in detail below may be used. Alternatively or additionally, the inventive compounds may be prepared using any of a variety combinatorial techniques, parallel synthesis and/or solid phase synthetic methods known in the art.

[0253] It will be appreciated as described below, that a variety of inventive compounds can be synthesized according to the methods described herein. The starting materials and reagents used in preparing these compounds are either available from commercial suppliers such as Aldrich Chemical Company (Milwaukee, WI), Bachem (Torrance, CA), Sigma (St. Louis, MO), or are prepared by methods well known to a person of ordinary skill in the art following procedures described in such references as Fieser and Fieser 1991, "Reagents for Organic Synthesis", vols 1-17, John Wiley and Sons, New York, NY, 1991; Rodd 1989 "Chemistry of Carbon Compounds", vols. 1-5 and supps, Elsevier Science Publishers, 1989; "Organic Reactions", vols 1-40, John Wiley and Sons, New York, NY, 1991; March 2001, "Advanced Organic Chemistry", 5th ed. John Wiley and Sons, New York, NY; and Larock 1989, "Comprehensive Organic Transformations", VCH Publishers. These schemes are merely illustrative of some methods by which the compounds of this invention can be synthesized, and various modifications to these schemes can be made and will be suggested to a person of ordinary skill in the art having regard to this disclosure.

[0254] The starting materials, intermediates, and compounds of this invention may be isolated and purified using conventional techniques, including

filtration, distillation, crystallization, chromatography, and the like. They may be characterized using conventional methods, including physical constants and spectral data.

[0255] General Reaction Procedures:

[0256] Unless mentioned specifically, reaction mixtures were stirred using a magnetically driven stirrer bar. An inert atmosphere refers to either dry argon or dry nitrogen. Reactions were monitored either by thin layer chromatography, by proton nuclear magnetic resonance (NMR) or by high-pressure liquid chromatography (HPLC), of a suitably worked up sample of the reaction mixture.

[0257] General Work Up Procedures:

[0258] Unless mentioned specifically, reaction mixtures were cooled to room temperature or below then quenched, when necessary, with either water or a saturated aqueous solution of ammonium chloride. Desired products were extracted by partitioning between water and a suitable water-immiscible solvent (e.g. ethyl acetate, dichloromethane, diethyl ether). The desired product containing extracts were washed appropriately with water followed by a saturated solution of brine. On occasions where the product containing extract was deemed to contain residual oxidants, the extract was washed with a 10% solution of sodium sulphite in saturated aqueous sodium bicarbonate solution, prior to the aforementioned washing procedure. On occasions where the product containing extract was deemed to contain residual acids, the extract was washed with saturated aqueous sodium bicarbonate solution, prior to the aforementioned washing procedure (except in those cases where the desired product itself had acidic character). On occasions where the product containing extract was deemed to contain residual bases, the extract was washed with 10% aqueous citric acid solution, prior to the aforementioned washing procedure (except in those cases where the desired product itself had basic character). Post washing, the desired product containing extracts were dried over anhydrous magnesium sulphate, and then filtered. The crude products were then isolated by removal of solvent(s) by rotary evaporation under reduced pressure, at an appropriate temperature (generally less than 45°C).

[0259] **General Purification Procedures:**

[0260] Unless mentioned specifically, chromatographic purification refers to flash column chromatography on silica, using a single solvent or mixed solvent as eluent. Suitably purified desired product containing elutes were combined and concentrated under reduced pressure at an appropriate temperature (generally less than 45°C) to constant mass. Final compounds were dissolved in 50% aqueous acetonitrile, filtered and transferred to vials, then freeze-dried under high vacuum before submission for biological testing.


[0261] ***Synthesis of Exemplary Compounds:***

[0262] The practitioner has a well-established literature of amino acid and peptide chemistry to draw upon, in combination with the information contained in the many examples which follow, for guidance on synthetic strategies, protecting groups, and other materials and methods useful for the synthesis of the compounds of this invention, including compounds containing the various R¹, R² and R³ substituents and X¹, X² and X³ moieties. Described generally below, are procedures and general guidance for the synthesis of compounds as described generally and in subclasses and species herein. In addition, synthetic guidance for the synthesis of amino acid derivatives and peptide analogues (and protease inhibitors more generally) can be found in U.S. Patent Nos: 5,585,397; 5,916,438 and 5,413,999; U.S. Application No. 10/462,127 filed June 16, 2003 and Published PCT application WO 02/02505, the entire contents of which are hereby incorporated by reference. A derivative of Formula I, or a pharmaceutically-acceptable derivative or salt thereof, may be prepared using any of the available relevant chemical transformations, combined with protection and deprotection as desired or required. Such processes, when used to prepare a derivative of the formula I, or a pharmaceutically-acceptable salt thereof, are illustrated by the following representative examples. The various starting materials are either commercially available or may be obtained by standard procedures of organic chemistry. The preparation of such starting materials is described within the accompanying non-limiting Examples.


[0263] Unless otherwise indicated, starting materials are either commercially available, as indicated or are obtained through laboratory synthesis by anyone reasonably familiar with the art.

[0264] EXAMPLE 6

[0265] This example describes the synthesis of amino acids and amino-acid like compounds of the formula




where P is a protecting group and R_A is substituted or unsubstituted aliphatic or aromatic moiety. These compounds were made as described in *J. Org. Chem.* **64**: 3322 (1999) and described by Scheme 1.

[0266] Scheme 1

[0267] Compound **1.2** was prepared by the treatment of (R,R)-(-)-pseudoephedrine (1 eq) and glycine methyl ester hydrochloride **1.1** (1.3 eq) in tetrahydrofuran (THF) with lithium tert-butoxide (1.4 eq) and worked up with water. Enolization of compound **1.2** in LiCl (3.2 eq) in THF by LHMDS (3.2 eq) and alkylation of $R_A X$ yields compound **1.3**. Hydrolysis of compound **1.3** under basic condition (NaOH, H_2O , refluxed) followed by protection of the amine gives compound **1.4**.


[0268] EXAMPLE 7

[0269] This example describes the synthesis of compounds of the formula



[0270] where R_A is as defined in Example 6. These compounds were prepared according to Scheme 2 and the procedure below.


[0271] Scheme 2



[0272] Compound 2.1 is prepared by the procedure described *J. Am. Chem. Soc.* 122: 3522 (2000) starting with the corresponding N-protected compound made by Example 6 or purchased from commercial sources. Reduction of lactone 2.1 by lithium aluminum hydride ("LAH") yields the corresponding diol 2.2, which was followed by the protection of the primary alcohol by tert-butyl-dimethyl silyl chloride ("TBSCl"; 1 eq), imidazole in CH₂Cl₂ to afford compound 2.3. Oxidation of the secondary alcohol by Dess-Martin periodinane to ketone 2.4, and reduction of ketone 2.4 by NaBH₄ afford alcohol 2.5. The mesylate 2.6 is obtained by the treatment of alcohol 2.5 with mesyl chloride. The addition of sodium azide to the mesylate 2.6 affords compound 1.

[0273] EXAMPLE 8


[0274] An alternate procedure to prepare compound 1 is described in Scheme 3 and the procedure below.

[0275] Scheme 3

[0276] Methyl ester **3.1** is prepared by esterification of the corresponding compound **1.4** (Example 6). The ketophosphonate **3.2** is made from methyl ester **3.1** by Claisen condensation with litho-dimethyl methylphosphonate. The Wadsworth-Emmons reaction of ketophosphonate **3.2** with ethyl pyruvate affords olefin **3.3**, which is followed by the reduction by NaBH_4 to afford lactone **3.4**. Hydrogenation of lactone **3.4** followed by reduction of LAH yields diol **3.6**. The protection of primary alcohol gives TBS ether **3.7**. The mesylation of **3.7** followed by the addition of sodium azide to the mesylate affords compound **1**.

[0277] EXAMPLE 9

[0278] Another alternate procedure to prepare compound **1** is described in Scheme 4 and the procedure below.


[0279] Scheme 4

[0280] Compound **4.2** and **4.3** are prepared by the procedure described *J. Org. Chem.* **56**: 4823 (1991), starting with the corresponding compound **1.4** (Example 6). Reduction of ketone **4.3** by NaBH_4 , followed by hydrogenation gives

the corresponding diol 4.4. The protection of the primary alcohol by TBSCl (1 eq), imidazole in CH_2Cl_2 affords compound 4.5. Mesylation of 4.5 followed by the addition of sodium azide to the mesylate 4.6 affords compound 1.


[0281] **EXAMPLE 10**


[0282] This example describes the synthesis of



[0283] which was prepared according to Scheme 5 and the procedure below.

[0284] **Scheme 5**





a) Compound 5.1 ([3-Methyl-1-(4-methyl-5-oxo-tetrahydro-furan-2-yl)-butyl]-carbamic acid tert-butyl ester) was prepared by the procedure described in *J. Am. Chem. Soc.* 122: 3522 (2000).

b) A solution of LiAlH₄ (40 mL, 1 M in THF) was added dropwise to 5.7 g of compound 5.1 in 40 ml of THF at 0 °C. The reaction solution was warmed to room temperature, stirred for 1.5 hours and cooled to 0 °C. A solution of 15% NaHSO₄ in water was added dropwise to the reaction mixture until no further precipitate was formed (6 ml of NaHSO₄). The reaction mixture was filtered, and concentrated, and the product 5.2 (5.4g, 93% yield) was used for next reaction without further purification. LCMS: 290(M+1).

c) TBSCl(2.82g, 18.8mmol) was added to the solution of diol 5.2 (5.4g, 18.8mmol) and imidazole (2.6g, 38.2 mmol) in 30 ml of CH₂Cl₂. After 30 minutes of stirring, the solvent was removed under reduced pressure. The reaction mixture was purified by column chromatography (40% ether / hexanes) to afford 6.5g (88% yield) of compound 5.3. LCMS: 404(M+1). ¹H NMR(CDCl₃)δ: 4.7(1H, broad s), 3.72(1H, m), 3.6-3.4(3H, m), 3.2(1H, broad s), 1.93 (1H, m), 1.7-1.55 (2H, m), 1.45(10H, m), 1.35-1.25 (2H, m), 0.95-0.85(18H, m), 0.04 (6H, s).

d) A solution of Dess-Martin periodate (5.5g, 13 mmol in 5.5 mL CH_2Cl_2) was added to a solution of compound **5.3** (3.5g, 8.68 mmol) in 30 mL of CH_2Cl_2 . After stirring for 15 minutes, the reaction mixture was purified by column (20% ether / hexanes) to afford compound **5.4** (3.2g, 91%). LCMS: 402(M+1).

e) To a solution of ketone **5.4** (3.2g, 8.0 mmol) in 50 mL of MeOH, NaBH_4 (0.3g) was added at -78°C . The reaction mixture was stirred for 10 minutes at -30°C , and was concentrated under reduced pressure. The reaction mixture was then extracted by ether (3 x 50 mL) water (50 mL) and washed with brine (50mL), and then concentrated to afford compound **5.5** (2.6g, 81% yield) as 20:1 ratio of syn and anti isomer (determined by ^1H NMR). LCMS: 404(M+1).

f) Mesyl chloride (1.1g, 9.7mmol) was added to the solution of alcohol **5.5** (2.6 g, 6.5mmol) and triethylamine (2 mL) in chloroform (20 mL) at 0°C . The resulting mixture was stirred for 60 minutes, then purified by column chromatography (20% ether / hexanes) to afford mesylate **5.6** (2.5g, 80%). LCMS: 482 (M+1).

g) Sodium azide (3.4g) was added to a solution of mesylate **5.6** (2.5g) in dimethylformamide ("DMF"; 20 mL). The reaction mixture was stirred for overnight at 75°C and extracted (ether 5x50mL/ Water 50 mL). The combined organic solution was dried, concentrated and purified by column chromatography (10%, ether/haxanes) to afford desired product **5.7** (1.05g, 55%), which used as common intermediate LCMS: 429(M+1) ^1H NMR(CDCl_3) δ : 4.43(1H, broad), 3.77(1H, m), 3.63 (1H, m), 3.49(2H, m), 1.85 (1H, m), 1.7-1.6 (2H, m), 1.45(10H, m), 1.35-1.25 (2H, m), 1.0-0.88(18H, m), 0.05(6H, s) and syn isomer (0.19g).

h) A solution of 4N HCl in dioxane was added to a solution of compound **5.7** in MeOH. After stirring at room temperature for 60 minutes, the volatiles were removed by reduced pressure to afford compound **5.8**, which was without further purification.

i) Method 1: To a mixture of compound 5.8 and N,N-dipropyl-isophthalamide in DMF, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride ("EDC")/1-hydroxybenzotriazole hydrate("HOBt")/diisopropylethylamine ("DIEA") was added. The resulting solution was stirred at room temperature overnight. 1N HCl (2 mL) was added, followed by addition of EtOAc (80 mL). After stirring for 10 minutes, the organic layer was separated, washed with brine and dried over MgSO₄. After removal of solvents, the residue was purified by column chromatography to afford compound 5.9. Method 2: To a mixture of compound 5.8 and N,N-dipropyl-isophthalamide in CH₂Cl₂, 2-(1H-azabenzotriazole-1-yl)-1,3,3-tetramethyluronium hexafluorophosphate("HATU")/DIEA was added. The resulting mixture was stirred at room temperature for 2 hours. The volatiles were removed by reduced pressure and purified by column chromatography to afford compound 5.9.


j) Compound 5.9 was oxidized by Jones' reagent, after worked up and purified by column to give acid 5.10. LCMS: 460(M+1) ¹H NMR(CD₃OD)δ: 7.92(1H, d), 7.80(1H, s), 7.5-7.6(2H, m), 4.33(1H, m), 3.59(1H, m), 3.51(2H, t), 3.22 (3H, t), 2.70 (1H, m), 1.98(1H, m), 1.80-1.35 (8H, m), 1.20(3H, t), 1.06-0.9(9H, m), 0.75(3H, t).

k) Method 1: To a mixture of compound 5.10 and 2-amino-N-benzyl-3-methylbutyramide in DMF, EDC/HOBt/ DIEA was added. The resulting solution was stirred at room temperature overnight. 1N HCl (2 mL) was added, followed by addition of EtOAc (80 mL). After stirring for 10 minutes, the organic layer was separated, washed with brine and dried over MgSO₄. After removal of solvents, the residue was purified by column chromatography to afford compound 5.11. Method 2: To a mixture of compound 5.10 and 2-amino-N-benzyl-3-methylbutyramide in CH₂Cl₂, HATU/DIEA was added. The resulting mixture was stirred at room temperature for 2 hours, the volatiles were removed by reduced pressure and purified by column chromatography to afford compound 5.11.

l) Method 1: Compound **5.11** is treated with 1M PhMe₃ in THF, and the resulting mixture is stirred for several hours until compound **5.11** is consumed. The reaction mixture is concentrated and 1N HCl is added. The resulting mixture is stirred overnight after which 20 % NaOH(aq) is added dropwise until the pH of the solution reaches 12-13. Method 2: Hydrogenation of compound **5.11** by 10 % Pd-C in MeOH yields compound **5.12** which is purified by preparative HPLC.


[0285] **EXAMPLE 11**


[0286] This example describes the synthesis of



which was prepared according to Scheme 6 and the procedure below.

[0303] **Scheme 6**





a) A solution of dimethyl methylphosphonate **6.1** (34.1 g, 215.4 mmol) in THF (250 mL) in a nitrogen atmosphere was cooled to -78°C and then was added 2.0 M solution of butyllithium (107 mL, 215.4 mmol) via canula in 20 minutes. The solution was stirred at -78°C for 20 minutes, and a THF (150 mL) solution of N-boc-L-phenylalanine methyl ester (10.0 g, 35.8 mmol) was slowly added via dropping funnel. The mixture was stirred at -78°C for 1 hour. The reaction was then quenched with 10% AcOH (250 mL) and warmed to room temperature. The solution was extracted with EtOAc, and the combined organic extracts were washed with saturated aqueous NaHCO_3 and brine and then dried over MgSO_4 . Solvents were evaporated and the excess dimethyl methylphosphonate was removed by rotary evaporator under high vacuum in a 90°C water bath. The resulted oily product **6.2** containing 5-10% dimethyl methylphosphonate (shown by ^1H NMR) was used without purification. ^1H NMR (400 MHz, CDCl_3) δ ppm 1.29 (m, 6 H) 1.35 (s, 9 H) 2.92 (dd, $J=14.24, 8.39$ Hz, 1 H) 3.03 (dd, $J=22.38, 13.73$ Hz, 1 H) 3.22 (m, 2 H) 4.10 (m, 4 H) 4.55 (d, $J=5.59$ Hz, 1 H) 5.38 (d, $J=7.88$ Hz, 1 H) 7.20 (m, 5 H); MS: 422 (MNa⁺)

b) A solution of the crude phosphonate **6.2** above (16.8 g, ca. 35.8 mmol) in THF (100 mL) in a nitrogen atmosphere was cooled to 0°C and then was added 1.6 M solution of butyllithium (22.4 mL, 35.8 mmol) via syringe in 10 minutes. The solution was stirred at 0°C for 30 minutes, and ethyl pyruvate (7.1 mL, 63.2 mmol) was added slowly. Stirring was continued at 0°C for 1 hour and at room temperature overnight. The reaction was quenched with saturated aqueous NH_4Cl (100 mL) and diluted with EtOAc (400 mL). The organic layer was separated,

washed with brine and then dried over MgSO_4 . Solvents were evaporated and the residue was purified by column chromatography (silica gel, Hexane-EtOAc 5:1) to give oily pure product **6.3** (9.0 g, 70% starting from **1**). ^1H NMR (400 MHz, ACETONE- D_6) δ ppm 1.24 (t, $J=7.12$ Hz, 3 H) 1.33 (s, 9 H) 2.00 (d, $J=1.27$ Hz, 3 H) 2.86 (m, 1 H) 3.19 (dd, $J=13.99, 5.09$ Hz, 1 H) 4.18 (q, $J=7.12$ Hz, 2 H) 4.53 (m, 1 H) 6.13 (d, $J=7.88$ Hz, 1 H) 6.51 (d, $J=1.27$ Hz, 1 H) 7.22 (m, 5 H); MS: 384 (MNa^+).

c) To a solution of compound **6.3** (8.4 g, 23.2 mmol) in MeOH (200 mL) cooled to -78°C was added sodium borohydride (1.7 g, 46.4 mmol). The mixture was stirred and warmed to -15°C in 3 hours, and then kept at the same temperature overnight. 1N aqueous HCl (100 mL) was added to the cold reaction mixture, and the volatiles were removed by rotary evaporator. The resulted mixture was extracted with EtOAc. The combined extracts were washed with saturated aqueous NaHCO_3 and brine and dried over MgSO_4 . After removal of solvents, the residue was purified by column chromatography (silica gel, Hexane-EtOAc 3:1) to give pure product **6.4** (4.3 g, 58%) and its 5-epimer (3.0 g, 41%) both as white solid. ^1H NMR (400 MHz, CDCl_3) δ ppm 1.36 (s, 9 H) 1.88 (s, 3 H) 2.80 (d, $J=6.61$ Hz, 2 H) 4.05 (m, 1 H) 4.68 (d, $J=8.39$ Hz, 1 H) 4.92 (s, 1 H) 6.91 (s, 1 H) 7.23 (m, 5 H); MS: 340 (Na^+).

d) Lactone **6.4** (3.5 g, 11.0 mmol) was dissolved in THF (300 mL), and to this solution 10% Pd/C (350 mg) was added. The mixture was stirred under an atmosphere of H_2 (balloon) for 5 hours, followed by filtration and concentration in vacuo, to provide product **6.5** (3.5 g, 99%) as white solid. ^1H NMR (400 MHz, CDCl_3) δ ppm 1.26 (d, $J=7.12$ Hz, 3 H) 1.35 (s, 9 H) 1.71 (m, 1 H) 2.42 (m, 1 H) 2.64 (m, 1 H) 2.87 (m, 1 H) 3.01 (m, 1 H) 3.92 (m, 1 H) 4.25 (m, 1 H) 4.46 (m, 1 H) 7.24 (m, 5 H); MS: 342 (MNa^+).

e) To a solution of compound **6.5** (3.5 g, 11.0 mmol) in THF (50 mL) cooled to 0°C was added 0.5 M LiAlH_4 in DME (22.0 mL, 11.0 mmol). The mixture was stirred at 0°C for 0.5 hours and warmed to room temperature. Stirring was continued until TLC indicated the reduction was completed. The mixture was

cooled to 0 °C and 1N NaHSO₄ (30 mL) was added slowly. After stirring for 30 minutes at room temperature, the mixture was extracted with EtOAc. The combined extracts were washed with saturated aqueous NaHCO₃ and brine and dried over MgSO₄. After removal of solvents, the residue was purified by column chromatography (silica gel, Hexane-EtOAc 1:1) to yield pure product **6.6** (2.7 g, 76%) as white solid. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.82 (d, J=6.61 Hz, 3 H) 1.19 (s, 9 H) 1.26 (m, 1 H) 1.41 (m, 1 H) 1.75 (m, J=15.13, 5.72 Hz, 1 H) 2.46 (dd, J=13.61, 10.05 Hz, 1 H) 2.93 (dd, J=13.73, 3.05 Hz, 1 H) 3.30 (m, 2 H) 3.50 (m, 2 H) 7.10 (m, 5 H); MS: 324 (MH⁺).

f) To a solution of compound **6.6** (1.0 g, 3.10 mmol) in DCM (15 mL) was added imidazole (422 mg, 6.20 mmol) and TBSCl (578 mg, 3.72 mmol) sequentially at 0 °C. After stirring at 0 °C for 1 hour, H₂O (10 mL) was added to quench the reaction. The organic layer was separated and the aqueous layer was extracted with DCM. The combined DCM solution was washed with H₂O and dried over MgSO₄. After removal of solvents, the residue was purified by column chromatography (silica gel, Hexane-EtOAc 3:1) to afford pure product **6.7** (1.34 g, 98%) as white solid. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.08 (s, 6 H) 0.87 (d, J=6.87 Hz, 3 H) 0.91 (s, 9 H) 1.32 (s, 9 H) 1.49 (m, J=6.36 Hz, 2 H) 1.80 (m, 1 H) 2.77 (m, 1 H) 2.89 (m, 1 H) 3.40 (m, 1 H) 3.58 (dd, J=10.05, 4.20 Hz, 1 H) 3.68 (s, 1 H) 3.81 (s, 1 H) 4.18 (m, J=12.33, 4.96 Hz, 1 H) 4.78 (d, J=8.65 Hz, 1 H) 7.21 (m, 5 H); MS: 438 (MH⁺), 460 (MNa⁺).

g) To a solution of compound **6.7** (1.2 g, 2.75 mmol) in chloroform (10 mL) was added triethylamine (0.8 mL, 5.5 mmol) and methanesulfonyl chloride ("MsCl"; 0.32 mL, 4.12 mmol) sequentially at 0 °C. After stirring at 0 °C for 1 hour, 1 N HCl (5 mL) was added to quench the reaction. The organic layer was separated and the aqueous layer was extracted with CH₂Cl₂. The combined CH₂Cl₂ solution was washed with H₂O and dried over MgSO₄. After removal of solvents, the residue was purified by column chromatography (silica gel, Hexane-EtOAc 3:1) to afford pure product **6.8** (1.34 g, 95%) as colorless oil. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.04 (s, 6 H) 0.88 (m, 12 H) 1.32 (s, 9 H) 1.37 (m, 1 H) 1.79 (m, 1 H) 1.95 (m, 1 H) 2.66

(dd, $J=13.22, 10.68$ Hz, 1 H) 2.90 (dd, $J=14.24, 4.83$ Hz, 1 H) 3.03 (s, 3 H) 3.40 (m, 1 H) 3.52 (m, 1 H) 4.16 (s, 1 H) 4.87 (d, $J=8.65$ Hz, 1 H) 4.96 (d, $J=9.16$ Hz, 1 H) 7.23 (m, 5 H); MS: 515 (MNa⁺).

h) Mesylate **6.8** (1.34 g, 2.60 mmol) was dissolved in anhydrous DMF (300 mL). To this stirred solution was added NaN₃ (2.50g, 39.0 mmol). The mixture was heated to 60 °C with an oil bath and stirred at this temperature for 8 hours. The solvent was evaporated in vacuo and then H₂O (20 mL) was added to dissolve the solid. The solution was extracted with ether, and the combined extracts were dried over MgSO₄. After removal of solvents, the residue was purified by column chromatography (silica gel, Hexane-EtOAc 10:1) to afford pure product **6.9** (0.72 g, 60%) as colorless oil. ¹H NMR (400 MHz, CDCl₃) δ ppm -0.05 (s, 3 H) -0.03 (s, 3 H) 0.74 (d, $J=6.87$ Hz, 3 H) 0.84 (s, 9 H) 1.39 (s, 9 H) 1.43 (m, 1 H) 1.63 (m, 1 H) 1.73 (m, 1 H) 2.75 (dd, $J=13.61, 8.77$ Hz, 1 H) 2.92 (m, 1 H) 3.37 (m, 2 H) 3.53 (t, $J=7.12$ Hz, 1 H) 3.91 (q, $J=8.22$ Hz, 1 H) 4.65 (d, $J=9.66$ Hz, 1 H) 7.25 (m, 5 H); MS: 485 (MNa⁺).

i) Azide **6.9** (109 mg, 0.24 mmol) was treated with 4N HCl in dioxane (5 mL). After stirring for 1 hour, the mixture was concentrated in vacuo and the resulted crude product **6.10** was used for the next step without purification. To a solution of the crude compound **6.10** in DMF (1 mL) was added N,N-dipropyl-isophthalamide acid (58 mg, 0.24 mmol), diisopropylethylamine (0.3 mL, 1.42 mmol), HOBT (54 mg, 0.35 mmol) and EDC (68 mg, 0.35 mmol) sequentially. The mixture was stirred at room temperature for 8 hours. 1N HCl (2 mL) was added, followed by addition of EtOAc (80 mL). After stirring for 10 minutes, the organic layer was separated, washed with brine and dried over MgSO₄. After removal of solvents, the residue was purified by column chromatography (silica gel, DCM-MeOH 19:1) to afford pure product **6.11** (113 mg, 81% starting from **6.9**) as colorless oil. ¹H NMR (400 MHz, CD₃OD) δ ppm 0.60 (t, $J=7.12$ Hz, 3 H) 0.81 (d, $J=6.61$ Hz, 3 H) 0.89 (t, $J=7.25$ Hz, 3 H) 1.32 (m, 1 H) 1.43 (m, 2 H) 1.64 (m, 3 H) 1.75 (m, 1 H) 2.84 (m, 1 H) 2.94 (m, 1 H) 3.06 (m, 2 H) 3.32 (m, 4 H) 3.61 (m, 1 H) 4.41 (m, 1 H) 7.06 (m, 1

H) 7.16 (q, J=7.46 Hz, 4 H) 7.40 (m, 2 H) 7.55 (s, 1 H) 7.67 (d, J=7.12 Hz, 1 H); MS: 480 (MH⁺).


j) To a solution of compound **6.11** (57 mg, 0.12 mmol) in acetone (1 mL) was added Jones' reagent (2.7M, 88 mM, 0.24 mmol). After stirring at room temperature for 20 minutes, iPrOH (40 mL) was added. Stirring was continued for 10 minutes and the reaction mixture was diluted with acetone (20 mL), followed by filtration over celite. The solution was dried (Na₂SO₄) and concentrated. The resulted crude product **6.12** was used for next step without purification.

k) To a DMF (0.5 mL) solution of the crude acid **6.12** was added 2-amino-N-benzyl-3-methyl-butyramide (20 mg, 0.075 mmol), diisopropylethylamine (0.1 mL, 0.45 mmol) and HATU (38 mg, 0.10 mmol) sequentially. The mixture was stirred at room temperature for 2 hours, and then purified by HPLC to provide pure product **6.13** (26 mg, 76% starting from compound **6.11**) as white solid. ¹H NMR (400 MHz, METHANOL-D) δ ppm 0.57 (t, J=7.25 Hz, 3 H) 0.76 (m, 6 H) 0.87 (t, J=7.25 Hz, 4 H) 0.98 (d, J=6.61 Hz, 3 H) 1.41 (m, 3 H) 1.59 (m, 2 H) 1.90 (m, 2 H) 2.66 (m, 1 H) 2.79 (dd, J=13.73, 9.66 Hz, 1 H) 2.94 (dd, J=13.99, 5.60 Hz, 1 H) 3.03 (m, 2 H) 3.22 (s, 1 H) 3.36 (m, 3 H) 3.99 (d, J=7.63 Hz, 1 H) 4.25 (s, 2 H) 4.40 (s, 1 H) 7.10 (m, 10 H) 7.38 (q, J=7.97 Hz, 2 H) 7.53 (s, 1 H) 7.65 (d, J=6.87 Hz, 1 H); MS: 682 (MH⁺).




l) To a solution of compound **6.13** (19 mg, 0.0279 mmol) in MeOH (5 mL) 10% Pd-C (6 mg) was added. The mixture was stirred under an atmosphere of H₂ for 30 minutes, followed by filtration and concentration. The resulting crude product was purified by preparative TLC to afford pure product **6.14** (16 mg, 88%) as white solid. ¹H NMR (400 MHz, METHANOL-D) δ ppm 0.61 (t, J=7.25 Hz, 3 H) 0.76 (d, J=6.61 Hz, 6 H) 0.88 (m, 4 H) 0.98 (d, J=6.87 Hz, 3 H) 1.28 (m, 1 H) 1.42 (m, 2 H) 1.62 (m, 2 H) 1.79 (m, 1 H) 1.91 (m, 1 H) 2.65 (dd, J=14.62, 6.23 Hz, 1 H) 2.75 (m, 2 H) 2.91 (m, 1 H) 3.07 (m, 2 H) 3.37 (m, 2 H) 4.00 (d, J=7.38 Hz, 1 H) 4.27 (m, 2 H) 7.05 (t, J=7.12 Hz, 1 H) 7.16 (m, 9 H) 7.40 (m, 2 H) 7.58 (s, 1 H) 7.71 (d, J=7.38 Hz, 1 H); MS: 656 (MH⁺).

[0287] EXAMPLE 12







[0288] The example describes the synthesis of compounds of the formula

























[0289] These compounds are prepared according the procedure of Example 10 or Example 11 except for using other amino acids and amino acid like compound

of the formula  as a reagent instead of  (Example 10) or  (Example 11). Illustrative examples of amino acids and amino acid like compounds and their corresponding products are shown in Table 1.

[0290] Table 1


Reagent	Product
	
	
	

Reagent	Product
	
	
	
	
	
	
	

Reagent	Product
	
	
	
	

[0291] **EXAMPLE 13**


[0292] This example describes the synthesis of



which was prepared according to Example 10 except for using 4-fluoroaniline instead of 2-amino-N-benzyl-3-methyl-butamide in step k. MS (M+1) 527.



[0293] **EXAMPLE 14**

[0294] The example describes the synthesis of compounds of the formula



[0295] These compounds are prepared according to Example 13 except for


using amino acids and amino acid-like compounds of the formula



instead of . Exemplary amino acids and amino acid-like compounds are shown in Table 1.

[0296] **EXAMPLE 15**


[0297] This example describes the synthesis of



which is prepared according to Example 10 except for using isopropylamine instead of 2-amino-N-benzyl-3-methyl-butylamide in step k when n=0 and for using isobutylamine instead of 2-amino-N-benzyl-3-methyl-butylamide in step k when n=1.



[0298] **EXAMPLE 16**

[0299] The example describes the synthesis of compounds of the formula




[0300] These compounds are prepared according to Example 15 except for


using amino acids and amino acid-like compounds of the formula




instead of .

[0301] EXAMPLE 17**[0302]** This example describes the synthesis of



which was prepared according to Example 10 except for using cyclopropylamine instead of 2-amino-N-benzyl-3-methyl-butylamide in step k when $n=0$ and cyclopropyl-methyl amine instead 2-amino-N-benzyl-3-methyl-butylamide in step k when $n=1$.

[0303] EXAMPLE 18**[0304]** The example describes the synthesis of compounds of the formula**[0305]** These compounds are prepared according to Example 17 except for


using amino acids and amino acid-like compounds of the formula




instead of

**[0306] EXAMPLE 19****[0307]** This example describes the synthesis of



which was prepared according to Example 10 except for 2-methyl-butylamine instead of 2-amino-N-benzyl-3-methyl-butylamide in step k.

[0308] EXAMPLE 20**[0309]** The example describes the synthesis of compounds of the formula**[0310]** These compounds are prepared according to Example 19 except for


using amino acids and amino acid-like compounds of the formula



instead of

**[0311] EXAMPLE 21****[0312]** This example describes the synthesis of







which is prepared according to Example 11 except for using 3-benzyloxy-2-oxo propanic acid methyl ester, which is prepared from 3-hydroxyl-2-oxo propanic acid instead of ethyl pyruvate in the step b, and 4-fluoroaniline instead of 2-amino-N-benzyl-3-methyl-butyramine in the step j.

[0313] EXAMPLE 22**[0314]** This example describes the synthesis of compounds of the formula

wherein R_B is a substituted or unsubstituted aliphatic or aromatic moiety. These compounds are prepared according to Example 21 except for using 2-oxo-acid esters of the formula $ROC(=O)C(=O)R_B$ (wherein R is lower alkyl and R_B is as defined


above) instead of 3-benzyloxy-2-oxo-propanic acid methyl ester. Exemplary esters and their corresponding products are shown in Table 2.

[0315] Table 2

Ester	Product
	
	
	













[0316] **EXAMPLE 23**

[0317] The example describes the synthesis of compounds of the formula




where R_C is substituted or unsubstituted aliphatic or aromatic moiety. These compounds are prepared according to Example 10 except for using acids of the formula $R_C\text{COOH}$ as a reagent instead of N,N-dipropyl-isophthalamic acid in step i and using 4-fluoroaniline instead of 2-amino-N-benzyl-3-methyl-butyramide in step k. Illustrative examples of acids and their corresponding products are shown in Table 3.

[0318] Table 3

R_cCOOH	Product
	
	
	
	
	
	


[0319] EXAMPLE 24

[0320] This example describes the synthesis of the compounds of the formula



which are prepared according to Scheme 7 and the procedure below.


[0321] Scheme 7




[0322] The Compounds 7.1 are prepared according to Example 16. These compounds are then reacted with phenyl chloroformate and DIEA to yield compounds 7.2. In certain embodiments, compounds 7.2 can be used as prodrug forms of their corresponding compounds 7.1.

[0323] EXAMPLE 25

[0324] This example describes the synthesis of compounds of the formula





















wherein R_D and R_E are each independently substituted or unsubstituted aliphatic or aromatic moieties. These compounds are prepared according to Scheme 8 and the procedure below.

[0325] Scheme 8

[0326] Compound 8.2 is treated with NaH in DMF, and halide R_DX (1 eq) is added to the reaction mixture to yield aryl ether 8.3. Compound 8.3 is then treated with NaH in DMF, and halide R_EX (1 eq) is added. The reaction mixture is hydrolyzed (LiOH, H_2O/THF) to give acid 8.1. Compounds 8.1, acids of the formula R_CCOOH , are used to make additional compounds of the invention. Table 4 shows exemplary compounds 8.1's.


[0327] Table 4

R_DX	R_EX	R_CCOOH
	CH_3I	

R_DX	R_EX	R_CCOOH
		
		
		
		
		
		


[0328] **EXAMPLE 26**

[0329] This example describes the synthesis of compounds of the formula



wherein R_F , R_G , and R_H are each independently substituted or unsubstituted aliphatic or aromatic. These compounds are prepared according to Scheme 9 and the procedure below.















[0330] Scheme 9























[0331] Mitsunobu reaction of compound 9.2 and the corresponding alcohol gives aryl ether 9.3. Hydrogenation of 9.3 reduces nitro group to aniline 9.4, which is followed by treatment of sulfonyl chloride to give the corresponding sulfonamide 9.5. Alkylation of sulfonamide 9.5 with halide (NaH , R_HX , 1 eq) gives 9.6, followed by basic hydrolysis ($LiOH$, $H_2O/THF/MeOH$) to give acid 9.1. Compounds 9.1, acids of the formula R_CCOOH , are used to make additional compounds of the invention. Table 5 shows exemplary compounds 9.1's.

[0332] Table 5

R_FX	R_GSO_2Cl	R_HX	R_CCOOH
	$MeSO_2Cl$	—	
	$MeSO_2Cl$	NT	


$R_F X$	$R_G SO_2 Cl$	$R_H X$	$R_C COOH$
	MeSO ₂ Cl		
CH ₃ I	MeSO ₂ Cl	CH ₃ I	
	MeSO ₂ Cl	CH ₃ I	
	MeSO ₂ Cl	CH ₃ I	
	MeSO ₂ Cl	CH ₃ I	
	MeSO ₂ Cl	CH ₃ I	
	MeSO ₂ Cl	CH ₃ I	

$R_F X$	$R_G SO_2 Cl$	$R_H X$	$R_C COOH$
	MeSO ₂ Cl	CH ₃ I	
	PhSO ₂ Cl	—I	
	PhSO ₂ Cl		
CH ₃ I	PhSO ₂ Cl	CH ₃ I	
	PhSO ₂ Cl	CH ₃ I	
	PhSO ₂ Cl	CH ₃ I	
	PhSO ₂ Cl	CH ₃ I	


$R_F X$	$R_G SO_2 Cl$	$R_H X$	$R_C COOH$
	$PhSO_2 Cl$	$CH_3 I$	
	$PhSO_2 Cl$	$CH_3 I$	
	$PhSO_2 Cl$	$CH_3 I$	

[0333] EXAMPLE 27

[0334] This example describes the synthesis of compounds of the formula


















wherein R_F is substituted or unsubstituted aliphatic or aromatic moiety. These compounds are prepared according to Scheme10 and the procedure below.

[0335] Scheme 10

[0336] Aniline 9.4 (Example 35) is treated with sulfonyl chloride to yield sulfonamide 10.2, which is followed by the treatment of NaH to afford cyclic sulfonamide 10.3. Basic hydrolysis (LiOH, H₂O/THF/MeOH) of compound 10.3 gives acid 10.1. Compounds 10.1, acids of the formula R_CCOOH, are used to make additional compounds of the invention. Table 6 shows exemplary compounds 10.1's.


[0337] Table 6

R _F	R _C COOH
	
	
—	


R_F	$R_C\text{COOH}$
	
	
	
	
	

[0338] **EXAMPLE 28**

[0339] This example describes the preparation of compounds of the formula



wherein R_I , R_J , and R_K are each independently substituted or unsubstituted aliphatic or aromatic moiety. These compounds are prepared according to Scheme 11 and the procedure below.




























[0340] Scheme 11




























[0341] The treatment of diamine **11.2** with sulfamine **a** under refluxing pyridine affords sulfamine **11.3**, which is followed by alkylation of α -halo alkyl ester to give compound **11.4**. Basic hydrolysis (LiOH, H₂O/THF/MeOH) of ester gives acid **11.1**. Compounds **11.1**, acids of the formula R_CCOOH, are used to make additional compounds of the invention. Table 7 shows exemplary compounds **11.1**'s.

[0342] Table 7

XCR ₁ R _J COOC	R _K NHCH ₂ CH ₂ N	R _C COOH
H ₃	H ₂	
Br-CH ₂ -CO ₂ Me		
Br-CH ₂ -CO ₂ Me		
Br-CH ₂ -CO ₂ Me		
Br-CH ₂ -CO ₂ Me		
Br-CH ₂ -CO ₂ Me		


$\text{XCR}_1\text{R}_1\text{COOC}$ H_3	$\text{R}_K\text{NHCH}_2\text{CH}_2\text{N}$ H_2	R_cCOOH

$\text{XCR}_1\text{R}_2\text{COOC}$ H_3	$\text{R}_3\text{NHCH}_2\text{CH}_2\text{N}$ H_2	R_4COOH
		
		
		
		
		
		
		
		
		


$\text{XCR}_1\text{R}_3\text{COOC}$	$\text{R}_K\text{NHCH}_2\text{CH}_2\text{N}$	R_cCOOH
H_3	H_2	
		
		
		
		
		
		
		
		
		

[0343] EXAMPLE 29

[0344] This example describes the synthesis of compounds of the formula



wherein R_I , R_J , and R_L are each independently substituted or unsubstituted aliphatic or aromatic moiety. These compounds are prepared according to Scheme 12 and the procedure below.

[0345] Scheme 12































[0346] Amino acid **12.2** is treated with benzyl amine and a coupling reagent such as HATU, to give **12.3**. Compound **12.3** is reduced by a reducing reagent such as LAH to give amine **12.4**. Compound **12.4** is treated with acid to afford diamine **12.5**, which follow by condensation with an oxalic acid derivative gives the dioxopiperazine **12.6**. Treatment of **12.6** with NaH and alkyl halide gives compound **12.7**. Removal of benzyl group by hydrogenation to give **12.8**, which is alkylated with an α -halo alkyl ester to give compound **12.9**. Hydrolysis of **12.9** affords acid **12.1**.













[0347] Compounds **12.1**, acids of the formula $R_C\text{COOH}$, are used to make additional compounds of the invention. Table 8 shows exemplary compounds **12.1**'s.

[0348] Table 8

$\text{XCR}_1\text{R}_2\text{COO}$ Me	R_1X	R_2COOH
$\text{BrCH}_2\text{CO}_2\text{Me}$	BuBr	
$\text{BrCH}_2\text{CO}_2\text{Me}$	PrBr	
$\text{BrCH}_2\text{CO}_2\text{Me}$		
$\text{BrCH}_2\text{CO}_2\text{Me}$		
$\text{BrCH}_2\text{CO}_2\text{Me}$		
$\text{BrCH}_2\text{CO}_2\text{Me}$		
$\text{BrCH}_2\text{CO}_2\text{Me}$		
$\text{BrCH}_2\text{CO}_2\text{Me}$		
$\text{BrCH}_2\text{CO}_2\text{Me}$	BuBr	


$\text{XCR}_1\text{R}_3\text{COO}$ Me	R_LX	R_cCOOH
	PrBr	
	BuBr	
	PrBr	

$\text{XCR}_1\text{R}_2\text{COO}$ Me	R_LX	R_cCOOH
		
		
		
		
		
		
		
		
		
		


$\text{XCR}_1\text{R}_2\text{COO}$ Me	R_LX	R_CCOOH
		
		
		
		

[0349] EXAMPLE 30

[0350] This example describes the synthesis of compounds of the formula















wherein R_M is substituted or unsubstituted aliphatic or aromatic moiety. These compounds are prepared according to Scheme 13 and the procedure below.

[0351] Scheme 13

[0352] Mitsunobu reaction of 13.2 gives the corresponding aryl ether 13.3, which is followed by the alkylation with halide (NaH , R_MX , 1 eq) to afford aryl diether 13.4. Basic hydrolysis (LiOH , $\text{H}_2\text{O}/\text{THF}/\text{MeOH}$) of 13.4 gives acid 13.1. Compounds 13.1, acids of the formula R_CCOOH , are used to make additional compounds of the invention. Table 9 shows exemplary compounds 13.1's.


[0353] Table 9

R_MX	R_CCOOH
	
	
	
	


R_MX	R_CCOOH
	
	

[0354] EXAMPLE 31

[0355] This example describes the synthesis of compounds of the formula

















wherein R_N , R_O , R_P , and R_Q are each independently substituted or unsubstituted aliphatic or aromatic moiety. These compounds are prepared according to Scheme 14 and the procedure below.















[0356] Scheme 14


[0357] The treatment of aniline **14.2** with sulfonyl chloride gives the corresponding sulfonamide **14.3**, which is followed by alkylation with halide to give **14.4**. The basic hydrolysis of **14.4** affords the mono-acid **14.5**, followed by amide coupling with corresponding amine to give **14.6**. Basic hydrolysis (LiOH,

H₂O/THF/MeOH) of 14.6 gives acid 14.1. Compounds 14.1, acids of the formula R_CCOOH, are used to make additional compounds of the invention. Table 10 shows exemplary compounds 14.1's.

[0358] Table 10


R _P NHR _Q	R _N SO ₂ Cl	R _O X	R _C COOH
	MeSO ₂ Cl	—I	
	MeSO ₂ Cl		
	PhSO ₂ Cl	—I	
	MeSO ₂ Cl	—I	
	MeSO ₂ Cl		
	PhSO ₂ Cl	—I	

$R_P NHR_Q$	$R_N SO_2 Cl$	$R_O X$	$R_C COOH$
	$MeSO_2 Cl$	$-I$	
	$MeSO_2 Cl$		
	$PhSO_2 Cl$	$-I$	
	$MeSO_2 Cl$	$-I$	
	$MeSO_2 Cl$		
	$PhSO_2 Cl$	$-I$	

[0359] EXAMPLE 32**[0360]** This example describes the synthesis of compounds of the formula

wherein R_R and R_S are each independently substituted or unsubstituted aliphatic or aromatic moiety, or R_R and R_S together form a substituted or unsubstituted cycloaliphatic or aromatic moiety. These compounds are prepared according to Scheme 15 and the procedure below.






















[0361] Scheme 15



















































[0362] Esterification of di-acid 15.2 with TMSCH_2N_2 gives the corresponding di-methyl ester 15.3. Basic hydrolysis (LiOH , 1 eq) of compound 15.3 yields mono-acid 15.4 which is then coupled with $\text{R}_\text{R}\text{NHR}_\text{S}$ to yield compound 15.5. The basic hydrolysis (LiOH , $\text{H}_2\text{O}/\text{THF}/\text{MeOH}$) of 15.5 gives acid 15.1. Compound 15.1, an acid of the formula $\text{R}_\text{C}\text{COOH}$, are used to make additional compounds of the invention. Table 11 shows exemplary compounds 15.1's.

























[0363] Table 11

























$R_R NHR_S$	$HO-C(=O)-Ar-C(=O)-OH$	$R_C COOH$






















R_RNHR_S	$HO-C(=O)-Ar-C(=O)-OH$	R_CCOOH
		
		
		
		
		
		
		

























$R_R NHR_S$	$HO-C(=O)-Ar-C(=O)-OH$	$R_C COOH$
		
		
		
		
		
		
		
		

























R_RNHR_S	$HO-C(=O)-\text{Ar}-C(=O)-OH$	R_CCOOH
		
		
		
		
		
		
		
		




























$R_R NHR_S$	$HO-C(=O)-Ar-C(=O)-OH$	$R_C COOH$
		
		
		
		
		
		
		
		




























R_RNHR_S	$HO-C(=O)-Ar-C(=O)-OH$	R_CCOOH
		
		
		
		
		
		
		
		




























R_RNHR_S	$HO-C(=O)-Ar-C(=O)-OH$	R_CCOOH
		
		
		
		
		
		
		

























$R_R NHR_S$	$HO-C(=O)-Ar-C(=O)-OH$	$R_C COOH$
		
		
		
		
		
		
		
		

$R_R NHR_S$	$HO-C(=O)-Ar-C(=O)-OH$	$R_C COOH$
		
		
		
		
		
		
		
		

R_RNHR_S	$HO-C(=O)-Ar-C(=O)-OH$	R_CCOOH
		
		
		
		
		
		
		
		
		

R_RNHR_S	$HO-C(=O)-Ar-C(=O)-OH$	R_CCOOH
		
		
		
		
		
		
		
		
		


R_RNHR_S	$HO-C(=O)-Ar-C(=O)-OH$	R_CCOOH
		
		
		
		
		
		
		
		
		

R_RNHR_S	$HO-C(=O)-\text{Ar}-C(=O)-OH$	R_CCOOH
		
		
		
		
		
		
		
		


R_RNHR_S	$HO-C(=O)-Ar-C(=O)-OH$	R_CCOOH

[0364] EXAMPLE 33

[0365] This example describes the synthesis of compounds of the formula




[0366] These compounds are prepared according to Example 23 except for using



[0367] as a reagent instead of

[0368] EXAMPLE 34


[0369] This example describes the synthesis of compounds of the formula




which are prepared according to Example 13 except for using acids of the formula $R_C\text{COOH}$ as a reagent instead of N,N-dipropyl-isophthalamide acid.

[0370] EXAMPLE 35

[0371] This example describes the synthesis of compounds of the formula




[0372] These compounds are prepared according to Example 34 except for using



[0373] EXAMPLE 36


[0374] This example describes the synthesis of





where R_T is substituted or unsubstituted aliphatic or aromatic moiety. These compounds are prepared according to Example 10 except for using amines of the formula H_2NR_T instead of 2-amino-N-benzyl-3-methyl-butylamide in step k.

[0375] EXAMPLE 37



[0376] This example describes the synthesis of compounds of the formula



[0377] These compound are prepared according to Example 36 except for using acids of the formula $R_C\text{COOH}$ as a reagent instead of N,N-dipropyl-isophthalamide acid.

[0378] EXAMPLE 38**[0379]** This example describes the synthesis of compounds of the formula**[0380]** These compounds are prepared according to Example 37 except for using

as a reagent instead of


[0381] EXAMPLE 39**[0382]** This example describes the synthesis of compounds of the formulawhere R_U is substituted or unsubstituted alkyl. These compounds are made according to Scheme 16 and the procedure below.**[0383] Scheme 16****[0384]** Compound 16.1 is prepared by amide coupling of compound 1.4 with N, O-dimethylhydroxylamine. Reaction of Weinreb amide 16.1 with Grignard reagent $R_U\text{MgX}$ yields ketone 16.2 which is then reduced to alcohol 16.3. Mesylation of compound 16.3 followed by the addition of azide replacement yields

compound **16.4** which is deprotected to yield amine **16.5**. Amide coupling of compound **16.5** and acid $R_C\text{COOH}$ with amide coupling reagents yields compound **16.6** which is reduced to make product **16.7**.

CLAIMS

What is claimed is:

1. An isolated compound having the structure:



or pharmaceutically acceptable derivative thereof;

wherein R' is hydrogen or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R' , taken together with R^2 or a substituent present on R^1 , may form a cycloheteroaliphatic moiety;

R^1 is an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^1 , taken together with R' , may form a cycloheteroaliphatic moiety;

X^1 is $-C(=O)-$, $-S(=O)-$, $-C(=NH)-$, $-C(=S)-$, $-NC(=O)-$, $-NC(=S)-$, $-N-C(=N-C\equiv N)-$, $-NS(O_2)-$, $-CHR^{X1A}-$, $-SO_2-$, $-COO-$, $-C(=O)C(R^{X1A})_2-$, or $-SC(=O)-$ wherein each occurrence of R^{X1A} is independently hydrogen, or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety;


R^2 is an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^2 , taken together with R' , may form a cycloheteroaliphatic moiety;

R^3 is hydrogen, halogen, or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety;

$R^{3'}$ is hydrogen, halogen, or lower alkyl;

R^4 is hydrogen, an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^4 , taken together with a substituent present on X^2 or X^3 , may form a cycloaliphatic, cycloheteroaliphatic, aromatic, or heteroaromatic moiety;

X^2 is absent, $-NR^{X2A}-$, $-(CHR^{X2A})_j-$, $-NR^{X2A}Y-$, $-(CHR^{X2A})_jY-$ or $-N(R^{X2A})CH(R^{X2A'})Y-$ wherein each occurrence of R^{X2A} is independently hydrogen or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety; each occurrence



of Y is independently $\left(\begin{array}{c} \text{O} \\ \parallel \\ \text{Y} \end{array} \right)_t$ wherein, for each independent occurrence of t , R^{X2B} is hydrogen, or an aliphatic, heteroaliphatic, aromatic or heteroaromatic

moiety, or R^{X2A} or one occurrence of R^{X2B} taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aromatic or heteroaromatic moiety, and wherein each occurrence of j and t is independently an integer from 1 to 4; and

X^3 is absent, $-NHCO-$, $-NHSO_2-$, $-NHCONH-$, $-NHCOO-$, $-CH_2NH-$, $-C(=O)-$, $-S(=O)-$, $-C(=NH)-$, $-C(=S)-$, $-NC(=S)N-$, $-N-C(=N-C\equiv N)N-$, $-NS(O_2)N-$, $-SO_2-$, $-C(=O)NR^{X3A}-$, $-C(=S)NR^{X3A}-$, $-COO-$, $-(CHR^{X3A})_k-$, $-O-$, $-CH_2NR^{X3A}-$, or $-NR^{X3A}-$, wherein each occurrence of R^{X3A} is independently hydrogen, an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^{X3A} taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aromatic or heteroaromatic moiety, and k is an integer from 1 to 3.

2. The compound of claim 1 wherein one or more of the following groups do not occur simultaneously as defined:

(i) R' , R^3 and $R^{3'}$ are each hydrogen; R^2 is alkyl, cycloalkylalkyl or aralkyl; $-X^2-X^3-R^4$ together represents $-CHR^eC(=O)NHCH(R^w)C(=O)NR^xR^y$, wherein R^e is hydrogen or alkyl, R^w is alkyl, and one of R^x or R^y represents hydrogen and the other represents hydrogen, alkyl, aryl, aralkyl, 1-alkoxycarbonyl-2-phenylethyl, 1-alkoxycarbonyl-2-(imidazol-4-yl)ethyl, 2-(imidazol-1-yl)ethyl, indanyl, heterocyclyl-alkyl, carboxyalkyl, alkoxycarbonylalkyl, aryloxyalkylalkyl, aralkoxycarbonylalkyl or a group of the formula $-A-N(R^a)(R^b)$ in which A represents alkylene and R^a and R^b each represents alkyl or R^a and R^b together represent a pentamethylene group in which one methylene group can be replaced by NH , N -alkyl, N -alkanoyl, N -aralkoxycarbonyl, O , S , SO , or SO_2 ; or R^x and R^y together with the nitrogen atom to which they are attached represent a 1,2,3,4-tetrahydroisoquinoline ring; and $-X^1-R^1$ together represents an alkoxycarbonyl, aralkoxycarbonyl, alkanoyl, aralkanoyl, aroyl, cycloalkylcarbonyl, heterocyclylcarbonyl, heterocyclyl-alkanoyl, 6-(dibenzylcarbamoyle)-4-oxohexanoyl moiety or an acyl group of an α -amino acid in which the amino group is substituted by an alkoxycarbonyl, aralkoxycarbonyl, diaralkylcarbamoyle, diaralkylalkanoyl, or aralkanoyl moiety;

wherein the term "aroyl" refers to an acyl group derived from an arylcarboxylic acid such as benzoyl, 1-naphthoyl, 2-naphthoyl, etc., and the term

“aralkanoyl” refers to an acyl group derived from an aryl-substituted alkanecarboxylic acid;

whereby the term “aryl” alone or in each of the aralkyl, aryloxycarbonylalkyl, aralkoxycarbonylalkyl or N-aralkoxycarbonyl moieties refers to a phenyl or naphthyl group optionally substituted with one or more substituents selected from alkyl, hydroxy, alkoxy and halogen;

(ii) R' , R^3 and $R^{3'}$ are each hydrogen; X^1 is $-C(=O)-$, $-SO_2-$, $N(R^x)SO_2$, $N(R^x)C(=O)$ or $SC(=O)$, wherein R^x is hydrogen, C_{1-5} alkyl or joined together with R^1 either directly to form a 5-7 membered heterocycle such as pyrrolidinyl or piperidinyl, or through a heteroatom selected from N, O and S, to form a 6-membered heterocycle with the nitrogen to which they are attached such as morpholinyl, piperazyl, or N- C_{1-3} alkyl-piperazyl; R^1 is a substituted or unsubstituted C_{1-6} alkyl, a 5-6 membered heterocycle or a 6-10 carbon atoms aryl moiety substituted with C_{1-4} alkyl, C_{1-3} alkoxy, hydroxy, halogen, $N(R^a)_2$, $C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-SO_2N(R^a)_2$, $CH_2N(R^a)_2$, $N(R^a)C(=O)R^a$ or $N(R^a)SO_2R^a$; R^2 is OR^a , $N(R^a)_2$, C_{1-4} alkenyl- R^c or $-[CR^bR^c]_nR^c$; and $-X^2-X^3-R^4$ together represent $-CH(R^d)C(=O)NHCH(R^e)C(=O)-Y-[CR^fR^g]_mR^g$, wherein m is an integer from 0 to 5, Y is O or NH, R^d and R^e are independently hydrogen, OR^a , $N(R^a)_2$, C_{1-4} alkenyl- R^c or $-[CR^bR^c]_nR^c$ wherein n is an integer from 0 to 5, R^a is hydrogen or C_{1-4} alkyl, R^b is hydrogen, hydroxy or C_{1-4} alkyl and R^c is hydrogen, substituted or unsubstituted aryl, 5- or 6-membered heterocycle, C_{1-6} alkyl or C_{1-6} alkenyl, C_{3-7} cycloalkyl, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring, benzofuryl, indolyl, azabicyclo C_{7-11} cycloalkyl or benzopiperidinyl; R^f is hydrogen, substituted or unsubstituted C_{1-6} alkyl or $(CH_2CH_2O)_pCH_3$ or $(CH_2CH_2O)_pH$ wherein p is an integer from 0 to 5, and R^g is hydrogen, or substituted or unsubstituted aryl, heterocycle, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring;

(iii) R' , R^3 and $R^{3'}$ are each hydrogen; X^1 is $-C(=O)-$, $-SO_2-$, $N(R^x)SO_2$, $N(R^x)C(=O)$ or $SC(=O)$, wherein R^x is hydrogen, C_{1-5} alkyl or joined together with R^1 either directly to form a 5-7 membered heterocycle such as pyrrolidinyl or piperidinyl, or through a heteroatom selected from N, O and S, to form a 6-membered heterocycle with the nitrogen to which they are attached such as

morpholinyl, piperazyl, or N-C₁₋₃alkyl-piperazyl; R¹ is a substituted or unsubstituted C₁₋₆alkyl, a 5-6 membered heterocycle or a 6-10 carbon atoms aryl moiety substituted with C₁₋₄alkyl, C₁₋₃alkoxy, hydroxy, halogen, N(R^a)₂, C(=O)OR^a, -C(=O)N(R^a)₂, -SO₂N(R^a)₂, CH₂N(R^a)₂, N(R^a)C(=O)R^a or N(R^a)SO₂R^a; R² is OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c; and -X²-X³-R⁴ together represent -CH(R^d)C(=O)-Y-[CR^fR^g]_mR^g, wherein m is an integer from 0 to 5, Y is O or NH, R^d is hydrogen, OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c; wherein n is an integer from 0 to 5, R^a is hydrogen or C₁₋₄alkyl, R^b is hydrogen, hydroxy or C₁₋₄alkyl and R^c is hydrogen, substituted or unsubstituted aryl, 5- or 6-membered heterocycle, C₁₋₆alkyl or C₁₋₆alkenyl, C₃₋₇cycloalkyl, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring, benzofuryl, indolyl, azabicyclo C₇₋₁₁cycloalkyl or benzopiperidinyl; R^f is hydrogen, substituted or unsubstituted C₁₋₆alkyl or (CH₂CH₂O)_pCH₃ or (CH₂CH₂O)_pH wherein p is an integer from 0 to 5, and R^g is hydrogen, or substituted or unsubstituted aryl, heterocycle, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring;

(iv) R¹ and R³ are each hydrogen; X¹ is -C(=O)-, -SO₂-, N(R^x)SO₂-, N(R^x)C(=O) or SC(=O), wherein R^x is hydrogen, C₁₋₅alkyl or joined together with R¹ either directly to form a 5-7 membered heterocycle such as pyrrolidinyl or piperidinyl, or through a heteroatom selected from N, O and S, to form a 6-membered heterocycle with the nitrogen to which they are attached such as morpholinyl, piperazyl, or N-C₁₋₃alkyl-piperazyl; R¹ is a substituted or unsubstituted C₁₋₆alkyl, a 5-6 membered heterocycle or a 6-10 carbon atoms aryl moiety substituted with C₁₋₄alkyl, C₁₋₃alkoxy, hydroxy, halogen, N(R^a)₂, C(=O)OR^a, -C(=O)N(R^a)₂, -SO₂N(R^a)₂, CH₂N(R^a)₂, N(R^a)C(=O)R^a or N(R^a)SO₂R^a; R² is OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c; R³ is hydrogen, OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c; and -X²-X³-R⁴ together represent -CH(R^d)C(=O)NHCH(R^e)C(=O)-Y-[CR^fR^g]_mR^g, wherein m is an integer from 0 to 5, Y is O or NH, R^d and R^e are independently hydrogen, OR^a, N(R^a)₂, C₁₋₄alkenyl-R^c or -[CR^bR^c]_nR^c wherein n is an integer from 0 to 5, R^a is hydrogen or C₁₋₄alkyl, R^b is hydrogen, hydroxy or C₁₋₄alkyl and R^c is hydrogen, substituted or unsubstituted aryl, 5- or 6-membered heterocycle, C₁₋₆alkyl or C₁₋₆alkenyl, C₃₋₇cycloalkyl, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring, benzofuryl, indolyl, azabicyclo C₇-


$_{11}$ cycloalkyl or benzopiperidinyl; R^f is hydrogen, substituted or unsubstituted C_{1-6} alkyl or $(CH_2CH_2O)_pCH_3$ or $(CH_2CH_2O)_pH$ wherein p is an integer from 0 to 5, and R^g is hydrogen, or substituted or unsubstituted aryl, heterocycle, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring;

(v) R' and $R^{3'}$ are each hydrogen; X^1 is $-C(=O)-$, $-SO_2-$, $N(R^x)SO_2$, $N(R^x)C(=O)$ or $SC(=O)$, wherein R^x is hydrogen, C_{1-5} alkyl or joined together with R^1 either directly to form a 5-7 membered heterocycle such as pyrrolidinyl or piperidinyl, or through a heteroatom selected from N, O and S, to form a 6-membered heterocycle with the nitrogen to which they are attached such as morpholinyl, piperazyl, or $N-C_{1-3}$ alkyl-piperazyl; R^1 is a substituted or unsubstituted C_{1-6} alkyl, a 5-6 membered heterocycle or a 6-10 carbon atoms aryl moiety substituted with C_{1-4} alkyl, C_{1-3} alkoxy, hydroxy, halogen, $N(R^a)_2$, $C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-SO_2N(R^a)_2$, $CH_2N(R^a)_2$, $N(R^a)C(=O)R^a$ or $N(R^a)SO_2R^a$; R^2 is OR^a , $N(R^a)_2$, C_{1-4} alkenyl- R^c or $-[CR^bR^c]_nR^c$; R^3 is hydrogen, OR^a , $N(R^a)_2$, C_{1-4} alkenyl- R^c or $-[CR^bR^c]_nR^c$; and $-X^2-X^3-R^4$ together represent $-CH(R^d)C(=O)-Y-[CR^fR^g]_mR^g$, wherein m is an integer from 0 to 5, Y is O or NH, R^d is hydrogen, OR^a , $N(R^a)_2$, C_{1-4} alkenyl- R^c or $-[CR^bR^c]_nR^c$; wherein n is an integer from 0 to 5, R^a is hydrogen or C_{1-4} alkyl, R^b is hydrogen, hydroxy or C_{1-4} alkyl and R^c is hydrogen, substituted or unsubstituted aryl, 5- or 6-membered heterocycle, C_{1-6} alkyl or C_{1-6} alkenyl, C_{3-7} cycloalkyl, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring, benzofuryl, indolyl, azabicyclo C_{7-11} cycloalkyl or benzopiperidinyl; R^f is hydrogen, substituted or unsubstituted C_{1-6} alkyl or $(CH_2CH_2O)_pCH_3$ or $(CH_2CH_2O)_pH$ wherein p is an integer from 0 to 5, and R^g is hydrogen, or substituted or unsubstituted aryl, heterocycle, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring; and

(vi) R' , R^3 and $R^{3'}$ are each hydrogen; R^2 is C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-7} cycloalkyl, aryl, heteroaryl, $T-C_{1-6}$ alkyl, $T-C_{2-6}$ alkenyl, wherein T is aryl, heteroaryl or C_{3-7} cycloalkyl; R^1-X^1 together represent W wherein W is R^x , R^xCO , R^xOCO , $R^xOCH(R^y)CO$, $R^xNHCH(R^y)CO$, $R^xSCH(R^y)CO$, R^xSO_2 , R^xSO or an amino acid with a blocked or unblocked amino terminus, wherein R^x and R^y are each independently hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, aryl, heteroaryl, $T-C_{1-6}$ alkyl or $T-(CH_2)_nCH(T)(CH_2)_n$ wherein n is an integer from 1 to 4; and $-X^2-X^3-R^4$ together

represent $-\text{CH}(\text{R}^a)\text{C}(=\text{X})\text{CHR}^b\text{R}^c$, wherein X is (OH,H) or O; R^a is hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-7} cycloalkyl, aryl, heteroaryl, T- C_{1-6} alkyl or T- C_{2-6} alkenyl; R^b is hydrogen or OH; and R^c is Y, $(\text{CHR}^w)_n\text{-Y}$ or $=\text{CR}^z(\text{CHR}^w)_n\text{-Y}$, wherein Y is hydrogen, OH, $-\text{NR}^q\text{R}^q$, aryl, heteroaryl or CO-Z, n is an integer from 1 to 4, Z is OH, $-\text{NR}^q\text{R}^q$, OR^w or an amino acid with a blocked or unblocked carboxy terminus, R^q is H, C_{1-6} alkyl or aryl C_{1-6} alkyl, and R^z and R^w are each independently hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, aryl, heteroaryl, T- C_{1-6} alkyl or T- C_{2-6} alkenyl.

3. The compound of claim 1 having the structure:



or pharmaceutically acceptable derivative thereof;


wherein R^1 is an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety;

R^2 is an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^2 , taken together with R^1 , may form a cycloheteroaliphatic moiety;

R^3 is hydrogen, halogen, or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety;

R^4 is hydrogen, an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^4 , taken together with a substituent present on X^2 or X^3 , may form a cycloaliphatic, cycloheteroaliphatic, aromatic, or heteroaromatic moiety;

X^2 is absent, $-\text{NR}^{\text{X}2\text{A}}-$, $-(\text{CHR}^{\text{X}2\text{A}})_j-$, $-\text{NR}^{\text{X}2\text{A}}\text{Y}-$, $-(\text{CHR}^{\text{X}2\text{A}})_j\text{Y}-$ or $-\text{N}(\text{R}^{\text{X}2\text{A}})\text{CH}(\text{R}^{\text{X}2\text{A}})\text{Y}-$ wherein each occurrence of $\text{R}^{\text{X}2\text{A}}$ is independently hydrogen or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, each occurrence



of Y is independently $\left(\begin{array}{c} \text{O} \\ \parallel \\ \text{R}^{\text{X}2\text{B}} \\ | \\ \text{N} \\ | \\ \text{H} \end{array} \right)_t$ wherein, for each independent occurrence of t, $\text{R}^{\text{X}2\text{B}}$ is hydrogen, or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or $\text{R}^{\text{X}2\text{A}}$ or one occurrence of $\text{R}^{\text{X}2\text{B}}$ taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aromatic or heteroaromatic moiety, and wherein each occurrence of j and t is independently an integer from 1 to 4; and

X^3 is absent, $-NHCO-$, $-NHSO_2-$, $-NHCONH-$, $-NHCOO-$, $-CH_2NH-$, $-C(=O)-$, $-S(=O)-$, $-C(=NH)-$, $-C(=S)-$, $-NC(=S)N-$, $-N-C(=N-C\equiv N)N-$, $-NS(O_2)N-$, $-SO_2-$, $-C(=O)NR^{X3A}-$, $-C(=S)NR^{X3A}-$, $-COO-$, $-(CHR^{X3A})_k-$, $-O-$, $-CH_2NR^{X3A}-$, or $-NR^{X3A}-$, wherein each occurrence of R^{X3A} is independently hydrogen, an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^{X3A} taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aromatic or heteroaromatic moiety, and k is an integer from 1 to 3.

4. The compound of claim 3 wherein one or more of the following groups do not occur simultaneously as defined:

(i) R^3 and $R^{3'}$ are each hydrogen; R^2 is alkyl, cycloalkylalkyl or aralkyl; $-X^2-R^4$ together represents $-CHR^eC(=O)NHCH(R^w)C(=O)NR^xR^y$, wherein R^e is hydrogen or alkyl, R^w is alkyl, and one of R^x or R^y represents hydrogen and the other represents hydrogen, alkyl, aryl, aralkyl, 1-alkoxycarbonyl-2-phenylethyl, 1-alkoxycarbonyl-2-(imidazol-4-yl)ethyl, 2-(imidazol-1-yl)ethyl, indanyl, heterocyclyl-alkyl, carboxyalkyl, alkoxycarbonylalkyl, aryloxycarbonylalkyl, aralkoxycarbonylalkyl or a group of the formula $-A-N(R^a)(R^b)$ in which A represents alkylene and R^a and R^b each represents alkyl or R^a and R^b together represent a pentamethylene group in which one methylene group can be replaced by NH , N -alkyl, N -alkanoyl, N -aralkoxycarbonyl, O , S , SO , or SO_2 ; or R^x and R^y together with the nitrogen atom to which they are attached represent a 1,2,3,4-tetrahydroisoquinoline ring; and $-C(=O)-R^1$ together represents an alkoxycarbonyl, aralkoxycarbonyl, alkanoyl, aralkanoyl, aroyl, cycloalkylcarbonyl, heterocyclylcarbonyl, heterocyclyl-alkanoyl, 6-(dibenzylcarbamoyl)-4-oxohexanoyl moiety or an acyl group of an α -amino acid in which the amino group is substituted by an alkoxycarbonyl, aralkoxycarbonyl, diaralkylcarbamoyl, diaralkylalkanoyl, or aralkanoyl moiety;

wherein the term "aroyl" refers to an acyl group derived from an arylcarboxylic acid such as benzoyl, 1-naphthoyl, 2-naphthoyl, etc., and the term "aralkanoyl" refers to an acyl group derived from an aryl-substituted alkanecarboxylic acid;

whereby the term "aryl" alone or in each of the aralkyl, aryloxy, carbonylalkyl, aralkoxycarbonylalkyl or N-aralkoxycarbonyl moieties refers to a phenyl or naphthyl group optionally substituted with one or more substituents selected from alkyl, hydroxy, alkoxy and halogen;

(ii) R^3 and $R^{3'}$ are each hydrogen; R^1 is a substituted or unsubstituted C_{1-6} alkyl, a 5-6 membered heterocycle or a 6-10 carbon atoms aryl moiety substituted with C_{1-4} alkyl, C_{1-3} alkoxy, hydroxy, halogen, $N(R^a)_2$, $C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-SO_2N(R^a)_2$, $CH_2N(R^a)_2$, $N(R^a)C(=O)R^a$ or $N(R^a)SO_2R^a$; R^2 is OR^a , $N(R^a)_2$, C_{1-4} alkenyl- R^c or $-[CR^bR^c]_nR^c$; and $-X^2-X^3-R^4$ together represent $-CH(R^d)C(=O)NHCH(R^e)C(=O)-Y-[CR^fR^g]_mR^g$, wherein m is an integer from 0 to 5, Y is O or NH, R^d and R^e are independently hydrogen, OR^a , $N(R^a)_2$, C_{1-4} alkenyl- R^c or $-[CR^bR^c]_nR^c$ wherein n is an integer from 0 to 5, R^a is hydrogen or C_{1-4} alkyl, R^b is hydrogen, hydroxy or C_{1-4} alkyl and R^c is hydrogen, substituted or unsubstituted aryl, 5- or 6-membered heterocycle, C_{1-6} alkyl or C_{1-6} alkenyl, C_{3-7} cycloalkyl, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring, benzofuryl, indolyl, azabicyclo C_{7-11} cycloalkyl or benzopiperidinyl; R^f is hydrogen, substituted or unsubstituted C_{1-6} alkyl or $(CH_2CH_2O)_pCH_3$ or $(CH_2CH_2O)_pH$ wherein p is an integer from 0 to 5, and R^g is hydrogen, or substituted or unsubstituted aryl, heterocycle, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring;

(iii) R^3 and $R^{3'}$ are each hydrogen; R^1 is a substituted or unsubstituted C_{1-6} alkyl, a 5-6 membered heterocycle or a 6-10 carbon atoms aryl moiety substituted with C_{1-4} alkyl, C_{1-3} alkoxy, hydroxy, halogen, $N(R^a)_2$, $C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-SO_2N(R^a)_2$, $CH_2N(R^a)_2$, $N(R^a)C(=O)R^a$ or $N(R^a)SO_2R^a$; R^2 is OR^a , $N(R^a)_2$, C_{1-4} alkenyl- R^c or $-[CR^bR^c]_nR^c$; and $-X^2-X^3-R^4$ together represent $-CH(R^d)C(=O)-Y-[CR^fR^g]_mR^g$, wherein m is an integer from 0 to 5, Y is O or NH, R^d is hydrogen, OR^a , $N(R^a)_2$, C_{1-4} alkenyl- R^c or $-[CR^bR^c]_nR^c$; wherein n is an integer from 0 to 5, R^a is hydrogen or C_{1-4} alkyl, R^b is hydrogen, hydroxy or C_{1-4} alkyl and R^c is hydrogen, substituted or unsubstituted aryl, 5- or 6-membered heterocycle, C_{1-6} alkyl or C_{1-6} alkenyl, C_{3-7} cycloalkyl, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring, benzofuryl, indolyl, azabicyclo C_{7-11} cycloalkyl or benzopiperidinyl; R^f is hydrogen, substituted or unsubstituted C_{1-6} alkyl or


$(\text{CH}_2\text{CH}_2\text{O})_p\text{CH}_3$ or $(\text{CH}_2\text{CH}_2\text{O})_p\text{H}$ wherein p is an integer from 0 to 5, and R^g is hydrogen, or substituted or unsubstituted aryl, heterocycle, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring;


(iv) $\text{R}^{3'}$ is hydrogen; R^1 is a substituted or unsubstituted C_{1-6} alkyl, a 5-6 membered heterocycle or a 6-10 carbon atoms aryl moiety substituted with C_{1-4} alkyl, C_{1-3} alkoxy, hydroxy, halogen, $\text{N}(\text{R}^a)_2$, $\text{C}(=\text{O})\text{OR}^a$, $-\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{SO}_2\text{N}(\text{R}^a)_2$, $\text{CH}_2\text{N}(\text{R}^a)_2$, $\text{N}(\text{R}^a)\text{C}(=\text{O})\text{R}^a$ or $\text{N}(\text{R}^a)\text{SO}_2\text{R}^a$; R^2 is OR^a , $\text{N}(\text{R}^a)_2$, C_{1-4} alkenyl- R^c or $-\text{[CR}^b\text{R}^c]_n\text{R}^c$; R^3 is hydrogen, OR^a , $\text{N}(\text{R}^a)_2$, C_{1-4} alkenyl- R^c or $-\text{[CR}^b\text{R}^c]_n\text{R}^c$; and $-\text{X}^2-\text{X}^3-\text{R}^4$ together represent $-\text{CH}(\text{R}^d)\text{C}(=\text{O})\text{NHCH}(\text{R}^e)\text{C}(=\text{O})-\text{Y}-\text{[CR}^f\text{R}^g]_m\text{R}^g$, wherein m is an integer from 0 to 5, Y is O or NH , R^d and R^e are independently hydrogen, OR^a , $\text{N}(\text{R}^a)_2$, C_{1-4} alkenyl- R^c or $-\text{[CR}^b\text{R}^c]_n\text{R}^c$ wherein n is an integer from 0 to 5, R^a is hydrogen or C_{1-4} alkyl, R^b is hydrogen, hydroxy or C_{1-4} alkyl and R^c is hydrogen, substituted or unsubstituted aryl, 5- or 6-membered heterocycle, C_{1-6} alkyl or C_{1-6} alkenyl, C_{3-7} cycloalkyl, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring, benzofuryl, indolyl, azabicyclo C_{7-11} cycloalkyl or benzopiperidinyl; R^f is hydrogen, substituted or unsubstituted C_{1-6} alkyl or $(\text{CH}_2\text{CH}_2\text{O})_p\text{CH}_3$ or $(\text{CH}_2\text{CH}_2\text{O})_p\text{H}$ wherein p is an integer from 0 to 5, and R^g is hydrogen, or substituted or unsubstituted aryl, heterocycle, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring;

(v) $\text{R}^{3'}$ is hydrogen; R^1 is a substituted or unsubstituted C_{1-6} alkyl, a 5-6 membered heterocycle or a 6-10 carbon atoms aryl moiety substituted with C_{1-4} alkyl, C_{1-3} alkoxy, hydroxy, halogen, $\text{N}(\text{R}^a)_2$, $\text{C}(=\text{O})\text{OR}^a$, $-\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{SO}_2\text{N}(\text{R}^a)_2$, $\text{CH}_2\text{N}(\text{R}^a)_2$, $\text{N}(\text{R}^a)\text{C}(=\text{O})\text{R}^a$ or $\text{N}(\text{R}^a)\text{SO}_2\text{R}^a$; R^2 is OR^a , $\text{N}(\text{R}^a)_2$, C_{1-4} alkenyl- R^c or $-\text{[CR}^b\text{R}^c]_n\text{R}^c$; R^3 is hydrogen, OR^a , $\text{N}(\text{R}^a)_2$, C_{1-4} alkenyl- R^c or $-\text{[CR}^b\text{R}^c]_n\text{R}^c$; and $-\text{X}^2-\text{X}^3-\text{R}^4$ together represent $-\text{CH}(\text{R}^d)\text{C}(=\text{O})-\text{Y}-\text{[CR}^f\text{R}^g]_m\text{R}^g$, wherein m is an integer from 0 to 5, Y is O or NH , R^d is hydrogen, OR^a , $\text{N}(\text{R}^a)_2$, C_{1-4} alkenyl- R^c or $-\text{[CR}^b\text{R}^c]_n\text{R}^c$; wherein n is an integer from 0 to 5, R^a is hydrogen or C_{1-4} alkyl, R^b is hydrogen, hydroxy or C_{1-4} alkyl and R^c is hydrogen, substituted or unsubstituted aryl, 5- or 6-membered heterocycle, C_{1-6} alkyl or C_{1-6} alkenyl, C_{3-7} cycloalkyl, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring, benzofuryl, indolyl, azabicyclo C_{7-11} cycloalkyl or benzopiperidinyl; R^f is hydrogen, substituted or unsubstituted C_{1-6} alkyl or

$(\text{CH}_2\text{CH}_2\text{O})_p\text{CH}_3$ or $(\text{CH}_2\text{CH}_2\text{O})_p\text{H}$ wherein p is an integer from 0 to 5, and R^8 is hydrogen, or substituted or unsubstituted aryl, heterocycle, 5- to 7-membered carbocyclic or 7- to 10-membered bicyclic carbocyclic ring; and

(vi) R^3 and $\text{R}^{3'}$ are each hydrogen; R^2 is C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-7} cycloalkyl, aryl, heteroaryl, T- C_{1-6} alkyl, T- C_{2-6} alkenyl, wherein T is aryl, heteroaryl or C_{3-7} cycloalkyl; $\text{R}^1\text{-C(=O)-}$ together represent W wherein W is R^x , R^xCO , R^xOCO , $\text{R}^x\text{OCH(R}^y\text{)CO}$, $\text{R}^x\text{NHCH(R}^y\text{)CO}$, $\text{R}^x\text{SCH(R}^y\text{)CO}$, R^xSO_2 , R^xSO or an amino acid with a blocked or unblocked amino terminus, wherein R^x and R^y are each independently hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, aryl, heteroaryl, T- C_{1-6} alkyl or T- $(\text{CH}_2)_n\text{CH(T)(CH}_2)_n$ wherein n is an integer from 1 to 4; and $-\text{X}^2\text{-X}^3\text{-R}^4$ together represent $-\text{CH(R}^a\text{)C(=X)CHR}^b\text{R}^c$, wherein X is (OH,H) or O; R^a is hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-7} cycloalkyl, aryl, heteroaryl, T- C_{1-6} alkyl or T- C_{2-6} alkenyl; R^b is hydrogen or OH; and R^c is Y, $(\text{CHR}^w)_n\text{-Y}$ or $=\text{CR}^z(\text{CHR}^w)_n\text{-Y}$, wherein Y is hydrogen, OH, $-\text{NR}^w\text{R}^q$, aryl, heteroaryl or CO-Z , n is an integer from 1 to 4, Z is OH, $-\text{NR}^w\text{R}^q$, OR^w or an amino acid with a blocked or unblocked carboxy terminus, R^q is H, C_{1-6} alkyl or aryl C_{1-6} alkyl, and R^z and R^w are each independently hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, aryl, heteroaryl, T- C_{1-6} alkyl or T- C_{2-6} alkenyl.

5. The compound of claim 3, wherein  has one of the following structures:




wherein R^3 is hydrogen, halogen or an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, $-(\text{heteroalkyl})\text{aryl}$, or $-(\text{heteroalkyl})\text{heteroaryl}$ moiety;

R^4 is an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, $-(\text{heteroalkyl})\text{aryl}$, or $-(\text{heteroalkyl})\text{heteroaryl}$ moiety, or R^4 , taken together with $\text{R}^{\text{X2A}'}$ or a substituent present on Y or X^3 , may form a cycloaliphatic, cycloheteroaliphatic, aryl, or heteroaryl moiety;


R^{X2A} is hydrogen or an aliphatic, heteroaliphatic, aryl, heteroaryl, - (alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety;

$R^{X2A'}$ is hydrogen, or an aliphatic, heteroaliphatic, aryl, heteroaryl, - (alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety, or $R^{X2A'}$ taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aryl or heteroaryl moiety;




Y is independently absent or is $\left(\begin{array}{c} \text{O} \\ \parallel \\ \text{---} \text{C} \text{---} \text{NH} \text{---} \text{C} \text{---} \text{R}^{X2B} \\ | \\ \text{H} \end{array} \right)_t$, wherein, for each independent occurrence of t , R^{X2B} is hydrogen, or an aliphatic, heteroaliphatic, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety, or one occurrence of R^{X2B} taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aryl or heteroaryl moiety; t is an integer from 1 to 4; and


X^3 is absent, -NHCO-, -NH SO_2 -, -NHCONH-, -NHCOO-, -CH $_2$ NH-, -C(=O)-, -S(=O)-, -C(=NH)-, -C(=S)-, -NC(=S)N-, -N-C(=N-C \equiv N)N-, -NS(O $_2$)N-, -SO $_2$ -, -C(=O)NR X3A -, -C(=S)NR X3A -, -COO-, (CHR X3A) $_k$, -O-, -CH $_2$ NR X3A -, or -NR X3A -, wherein each occurrence of R^{X3A} is independently hydrogen, an aliphatic, heteroaliphatic, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety, or R^{X3A} taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aryl or heteroaryl moiety, and k is an integer from 1 to 3.




6. The compound of claim 5, wherein $\begin{array}{c} \text{X}^2 \text{---} \text{X}^3 \text{---} \text{R}^4 \\ | \quad | \\ \text{R}^3 \quad \text{R}^{3'} \end{array}$ has one of the following structures:




wherein R^3 , R^4 , R^{X2A} , $R^{X2A'}$, R^{X2B} and t are as defined in claim 2.

7. The compound of claim 5, wherein  has one of the following structures:



wherein R^3 and R^4 are as defined in claim 2.

8. The compound of claim 3 having the structure:




wherein R^{X2A} is hydrogen or an aliphatic, heteroaliphatic, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety.

9. The compound of claim 8, wherein R^{X2A} is a substituted or unsubstituted, linear or branched lower alkyl moiety.


10. The compound of claim 5, wherein R^{X2A} is methyl, ethyl, propyl, isopropyl or phenethyl.

11. The compound of claim 3 or 8, wherein R^1 has one of the following structures:




wherein R^{1A} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, $-\text{OR}^{1B}$, $-\text{SR}^{1B}$, $-\text{N}(\text{R}^{1B})_2$, $-\text{SO}_2\text{N}(\text{R}^{1B})_2$, $-\text{C}(=\text{O})\text{N}(\text{R}^{1B})_2$, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OR}^{1B}$, $\text{N}(\text{R}^{1B})\text{C}(=\text{O})\text{R}^{1C}$ or $-\text{N}(\text{R}^{1B})\text{SO}_2\text{R}^{1C}$; wherein each occurrence of R^{1B} and R^{1C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$; or R^{1B} and R^{1C} , taken together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclic moiety; R^{1D} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$, $-(\text{alkyl})\text{heteroaryl}$, acyl or a nitrogen protecting group; wherein n and p are each independently integers from 0 to 3 and r is an integer from 1 to 6; whereby each of the foregoing alkyl and heteroalkyl moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, and each of the foregoing aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ and $-(\text{alkyl})\text{heteroaryl}$ moieties may be substituted or unsubstituted.

12. The compound of claim 11, wherein R^1 has one of the following structures:




13. The compound of claim 11, wherein R^1 has one of the following structures:




wherein R^{1D} is hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl; -(alkyl)heteroaryl or acyl; R^{1E} and R^{1F} are each independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -C(=O) R^{1C} or -SO₂ R^{1C} , where R^{1C} is hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl; or R^{1E} and R^{1F} taken together form a 5-8 membered heterocyclic ring; or R^{1E} and one occurrence of R^{1A} , taken together, form a substituted or unsubstituted, saturated or unsaturated heterocyclic ring.


14. The compound of claim 11, wherein R^1 has one of the following structures:



15. The compound of claim 11, wherein R^1 has one of the following structures:




16. The compound of claim 11, wherein R^1 has one of the following structures:



17. The compound of claim 3 or 8, wherein R^2 is lower alkyl, $-\text{CH}_2\text{NR}^{2A}\text{R}^{2B}$ or $-(\text{CH}_2)\text{phenyl}$, wherein the phenyl group is optionally substituted with one or more occurrences of R^{2C} , wherein R^{2C} is hydrogen, alkyl, alkoxy or halogen; and wherein R^{2A} and R^{2B} are each independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(\text{alkyl})\text{aryl}$ or $-(\text{alkyl})\text{heteroaryl}$; whereby each of the foregoing alkyl and heteroalkyl moieties may be linear or branched, substituted or unsubstituted,


cyclic or acyclic, and each of the foregoing aryl, heteroaryl, -(alkyl)aryl and -(alkyl)heteroaryl moieties may be substituted or unsubstituted.

18. The compound of claim 17, wherein R^2 has one of the following structures:



wherein R^{2A} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl, $-OR^{2B}$, $-SR^{2B}$, $-N(R^{2B})_2$, $-SO_2N(R^{2B})_2$, $-C(=O)N(R^{2B})_2$, halogen, $-CN$, $-NO_2$, $-C(=O)OR^{2B}$, $-N(R^{2B})C(=O)R^{2C}$, wherein each occurrence of R^{2B} and R^{2C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl, wherein q and s are each independently integers from 0 to 3 and u is an integer from 1 to 6; whereby each of the foregoing alkyl and heteroalkyl moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, and each of the foregoing aryl, heteroaryl, -(alkyl)aryl and -(alkyl)heteroaryl moieties may be substituted or unsubstituted.


19. The compound of claim 17, wherein R^2 has one of the following structures:




wherein each occurrence of R^{2A} is independently hydrogen or lower alkyl; each occurrence of X is independently a halogen; s is an integer from 0 to 3 and u is an integer from 1 to 6; whereby each of the foregoing alkyl moieties may be linear or branched, substituted or unsubstituted and cyclic or acyclic.

20. The compound of claim 17, wherein each occurrence of X is independently chlorine or fluorine and R^{2A} is methyl.


21. The compound of claim 17, wherein R^2 has one of the following structures:



22. The compound of claim 17, wherein R^2 has one of the following structures:




23. The compound of claim 3, wherein R^3 is hydrogen or halogen.
24. The compound of claim 3 or 8, wherein R^4 is substituted or unsubstituted, linear or branched, cyclic or acyclic alkyl, phenyl or $-(CH_2)phenyl$, wherein the phenyl group is optionally substituted with one or more occurrences of R^{4A} , wherein R^{4A} is hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, $-(alkyl)aryl$ or $-(alkyl)heteroaryl$, $-OR^{4B}$, $-SR^{4B}$, $-N(R^{4B})_2$, $-SO_2N(R^{4B})_2$, $-C(=O)N(R^{4B})_2$, halogen, $-CN$, $-NO_2$, $-C(=O)OR^{4B}$, $-N(R^{4B})C(=O)R^{4C}$, wherein each occurrence of R^{4B} and R^{4C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(alkyl)aryl$, $-(alkyl)heteroaryl$, $-(heteroalkyl)aryl$, $-(heteroalkyl)heteroaryl$.
25. The compound of claim 24, wherein R^4 is substituted or unsubstituted, linear or branched, cyclic or acyclic alkyl, phenyl or $-(CH_2)phenyl$, wherein the phenyl group is optionally substituted with one or more occurrences of R^{4A} , wherein R^{4A} is hydrogen, hydroxyl, alkyl, alkoxy or halogen.
26. The compound of claim 24, wherein R^4 has one of the following structures:



wherein each occurrence of R^{4A} is independently hydrogen, alkyl, heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl, $-OR^{4B}$, $-SR^{4B}$, $-N(R^{4B})_2$, $-SO_2N(R^{4B})_2$, $-C(=O)N(R^{4B})_2$, halogen, $-CN$, $-NO_2$, $-C(=O)OR^{4B}$, $-N(R^{4B})C(=O)R^{4C}$, wherein each occurrence of R^{4B} and R^{4C} is independently hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, -(alkyl)aryl or -(alkyl)heteroaryl, wherein v and w are each independently integers from 0 to 3 and x is an integer from 1 to 6; whereby each of the foregoing alkyl and heteroalkyl moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, and each of the foregoing aryl, heteroaryl, -(alkyl)aryl and -(alkyl)heteroaryl moieties may be substituted or unsubstituted.


27. The compound of claim 24, wherein R^4 is methyl, ethyl, propyl or one of:



wherein each occurrence of R^{4A} is independently hydrogen, lower alkyl or $C(=O)OR^{4B}$, wherein R^{4B} is hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(alkyl)aryl$ or $-(alkyl)heteroaryl$; each occurrence of X is independently a halogen; w is an integer from 0 to 3 and x is an integer from 1 to 6; whereby each of the foregoing alkyl and heteroalkyl moieties may be linear or branched, substituted or unsubstituted, cyclic or acyclic, and each of the foregoing aryl, heteroaryl, $-(alkyl)aryl$ and $-(alkyl)heteroaryl$ moieties may be substituted or unsubstituted.


28. The compound of claim 27, wherein each occurrence of X is independently chlorine or fluorine and R^{4A} is methyl.

29. The compound of claim 24, wherein R^4 is methyl, ethyl, propyl or one of:




wherein R^{4A} is hydrogen, hydroxyl, lower alkyl, lower alkoxy, halogen, $C(=O)OR^{4B}$, aryl, heteroaryl, $-(alkyl)aryl$ or $-(alkyl)heteroaryl$, wherein R^{4B} is hydrogen, lower alkyl, lower heteroalkyl, aryl, heteroaryl, $-(alkyl)aryl$, $-(alkyl)heteroaryl$, $-(heteroalkyl)aryl$ or $-(heteroalkyl)heteroaryl$.

30. The compound of claim 24, wherein R^4 is methyl, ethyl, propyl, isopropyl or one of:



31. The compound of claim 24, wherein R⁴ has one of the following structures:



32. A pharmaceutical composition comprising a compound of claim 3 or 8, and a pharmaceutically acceptable carrier or diluent, optionally further comprising an additional therapeutic agent.

33. The pharmaceutical composition of claim 32, wherein the compound is present in an amount effective to inhibit β -secretase activity.

34. The pharmaceutical composition of claim 32, wherein the additional therapeutic agent is an agent for the treatment of Alzheimer's Disease.

35. A method for inhibiting β -secretase activity in a patient or a biological sample, comprising administering to said patient, or contacting said biological sample with an effective inhibitory amount of a compound of claim 3 or 8.

36. A method for treating or preventing a disease characterized by β -amyloid deposits in the brain comprising administering to a patient a therapeutically effective amount of a compound of claim 3 or 8.


37. The method of claim 36, wherein the disease is Alzheimer's disease MCI (mild cognitive impairment), Down's syndrome, Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch-Type, cerebral amyloid angiopathy, other degenerative dementias, including dementias of mixed vascular and degenerative origin, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear palsy, dementia associated with cortical basal degeneration, and diffuse Lewy body type Alzheimer's disease.

AMENDED CLAIMS

WO 2005/005374 received by the International Bureau on 18 June 2004 (18.06.0PCT/US2003/039276
original claims 3, 5, 6, 7, 10 and 20 replaced by new claims 3, 5,
6, 7, 10 and 20; and 1-2, and 4 cancelled; remaining
claims unchanged (8 pages)]

+ STATEMENT

3. An isolated compound having the structure:



or pharmaceutically acceptable derivative thereof;


wherein R^1 and R^2 are independently an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety; or R^2 , taken together with R^1 , may form a cycloheteroaliphatic moiety;

R^3 is hydrogen, halogen, or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety;

$R^{3'}$ is hydrogen, halogen, or lower alkyl;

R^4 is hydrogen, an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^4 , taken together with a substituent present on X^2 or X^3 , may form a cycloaliphatic, cycloheteroaliphatic, aromatic, or heteroaromatic moiety;

X^2 is absent, $-NR^{X2A}-$, $-(CHR^{X2A})_j-$, $-NR^{X2A}Y-$, $-(CHR^{X2A})_jY-$ or $-N(R^{X2A})CH(R^{X2A'})Y-$ wherein each occurrence of R^{X2A} is independently hydrogen or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, each occurrence




of Y is independently $\left(\begin{array}{c} \text{O} \\ \parallel \\ \text{---} \text{N} \text{---} \text{H} \end{array} \right)_t$ wherein, for each independent occurrence of t , R^{X2B} is hydrogen, or an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^{X2A} or one occurrence of R^{X2B} taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aromatic or heteroaromatic moiety, and wherein each occurrence of j and t is independently an integer from 1 to 4; and

X^3 is absent, $-NHCO-$, $-NHSO_2-$, $-NHCONH-$, $-NHCOO-$, $-CH_2NH-$, $-C(=O)-$, $-S(=O)-$, $-C(=NH)-$, $-C(=S)-$, $-N(R^{X3A})C(=S)N(R^{X3A})-$, $-N(R^{X3A})-C(=N-C\equiv N)N(R^{X3A})-$, $-N(R^{X3A})S(O_2)N(R^{X3A})-$, $-SO_2-$, $-C(=O)NR^{X3A}-$, $-C(=S)NR^{X3A}-$, $-COO-$, $-(CHR^{X3A})_k-$, $-O-$, $-CH_2NR^{X3A}-$, or $-NR^{X3A}-$, wherein each occurrence of R^{X3A} is independently hydrogen, an aliphatic, heteroaliphatic, aromatic or heteroaromatic moiety, or R^{X3A} taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aromatic or heteroaromatic moiety, and k is an integer from 1 to 3;


AMENDED SHEET (ARTICLE 19)

with the proviso that

(i) the compound does not have one of the structures:



(ii) when $-C(R^3)(R^3)-X^2-X^3-R^4$ has the structure:



wherein R^{X2A} is hydrogen or C_{1-8} alkyl;

R^{X3A} is C_{1-8} alkyl;

one of R^{4A} or R^{4B} represents hydrogen and the other represents hydrogen, C_{1-8} alkyl, Ar, $-C_{1-8}$ alkyl-Ar, 1- C_{1-8} alkoxycarbonyl-2-phenylethyl, 1- C_{1-8} alkoxycarbonyl-2-(imidazol-4-yl)ethyl, 2-(imidazol-1-yl)ethyl, indanyl, heterocyclyl- C_{1-8} alkyl-, carboxy C_{1-8} alkyl, $-C_{1-8}$ alkylC(=O)OC $_{1-8}$ alkyl, $-C_{1-8}$ alkylC(=O)OAr, $-C_{1-8}$ alkylC(=O)OC $_{1-8}$ alkylAr or a group of the formula $-A-N(R^a)(R^b)$ in which A represents C_{1-8} alkylene and R^a and R^b each represents C_{1-8} alkyl or R^a and R^b together represent a pentamethylene group in which one methylene group can be replaced by NH, N- C_{1-8} alkyl, N- C_{1-8} alkanoyl, N-aralkoxycarbonyl, O, S, SO, or SO₂; or R^{4A} and R^{4B}


AMENDED SHEET (ARTICLE 19)

together with the nitrogen atom to which they are attached represent a 1,2,3,4-tetrahydroisoquinoline ring;

then R^1 is not $-OC_{1-8}alkyl$, $-OC_{1-8}alkylAr$, $-C_{1-8}alkyl$, $-C_{1-8}alkylAr$, $-OAr$, $-C_{3-8}cycloalkyl$, Het, $-C_{1-8}alkyl-Het$, $(PhCH_2)_2NC(=O)CH_2CH_2C(=O)CH_2CH_2-$ or the aminoalkyl radical of an α -amino acid residue in which the amino group is substituted by $-C(=O)OC_{1-8}alkyl$, $-C(=O)OC_{1-8}alkylAr$, $-C(=O)N(C_{1-8}alkylAr)_2$, $-C(=O)C_{1-8}alkyl(C_{1-8}alkylAr)_2$ or $-C(=O)C_{1-8}alkylAr$;

wherein Ar is phenyl or naphthyl optionally substituted with one or more substituents selected from halogen, $-OH$, $-C_{1-8}alkoxy$ and $C_{1-8}alkyl$; and Het is a saturated, partially unsaturated or aromatic monocyclic, bicyclic or tricyclic heterocycle which is attached via a carbon atom, contains one or more heteroatoms selected from N, O and S, and is optionally substituted on one or more carbon atoms by $C_{1-8}alkyl$, $C_{1-8}alkoxy$ and/or halogen; and

(iii) when $-C(R^3)(R^3)-X^2-X^3-R^4$ has the structure:



wherein each occurrence of n is independently 0 or 1;


R^{X2B} is hydrogen, $C_{1-6}alkyl$; $-CH_2Ar$, cyclohexyl or adamantyl;


W is $-OR^{W1}$; $-N(R^{W1})_2$; $-NR^{W2}-(CH_2)_m-Q_1$; an optionally amide- or ester-protected serine, lysine or arginine residue, or optionally ester- or ether-protected amino alcohol derivative thereof; or an N-containing heterocyclic ring attached via the N atom containing 1-3 heteroatoms selected from N, O and S, optionally substituted with one or more substituents selected from R^{W1} and $-CH_2R^{W1}$; wherein m is 2-6; Q_1 is NH_2 or $-NH-C(=NH)NH_2$ wherein any of the hydrogen atoms attached to the nitrogen atoms may be substituted with $C_{1-6}alkyl$, cycloalkyl or R^{W2} ; R^{W1} , for each occurrence, is independently hydrogen, $C_{1-6}alkyl$, $C_{3-7}cycloalkyl$, $C_{7-12}bi$ or tricycloalkyl, Ar or $-C_{1-6}alkyl-Ar$; and R^{W2} is hydrogen, $-(CH_2)_p-R^{W3}$, $-C(=O)-R^{W3}$, $-COOR^{W3}$, $-C(=O)-(CH_2)_pR^{W3}$, $-C(=O)O(CH_2)_pR^{W3}$, $-C(=O)-(CH_2)_pOR^{W3}$, $-SO_2-R^{W3}$ or -

$C(=O)N(R^{W3})_2$, wherein p is 0-5; and R^{W3} for each occurrence takes the definition of R^{W1} above;

then R^1 is not hydrogen, $-R^{1A}$, $-OR^{1A}$, $-(CH_2)_pR^{1A}$, $-O(CH_2)_pR^{1A}$, $-(CH_2)_pOR^{1A}$ or $-N(R^{1A})_2$, wherein p is 0-5; and R^{1A} for each occurrence takes the definition of R^{W1} above;

wherein Ar is phenyl or naphthyl optionally substituted with one or more substituents selected from halogen, $-OH$, $C_{1-6}alkyl$ and $-CF_3$.

5. The compound of claim 3, wherein  has one of the following structures:




wherein R^3 is hydrogen, halogen or an aliphatic, heteroaliphatic, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety;

R^4 is an aliphatic, heteroaliphatic, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety, or R^4 , taken together with $R^{X2A'}$ or a substituent present on Y or X^3 , may form a cycloaliphatic, cycloheteroaliphatic, aryl, or heteroaryl moiety;

R^{X2A} is hydrogen or an aliphatic, heteroaliphatic, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety;


$R^{X2A'}$ is hydrogen, or an aliphatic, heteroaliphatic, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety, or $R^{X2A'}$ taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aryl or heteroaryl moiety;


Y is independently absent or is , wherein, for each independent occurrence of t, R^{X2B} is hydrogen, or an aliphatic, heteroaliphatic, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety, or one occurrence of R^{X2B} taken together with R^4

AMENDED SHEET (ARTICLE 19)


may form a cycloaliphatic, cycloheteroaliphatic, aryl or heteroaryl moiety; t is an integer from 1 to 4; and


X^3 is absent, $-NHCO-$, $-NHSO_2-$, $-NHCONH-$, $-NHCOO-$, $-CH_2NH-$, $-C(=O)-$, $-S(=O)-$, $-C(=NH)-$, $-C(=S)-$, $-N(R^{X3A})C(=S)N(R^{X3A})-$, $-N(R^{X3A})-C(=N-C\equiv N)N(R^{X3A})-$, $-N(R^{X3A})S(O_2)N(R^{X3A})-$, $-SO_2-$, $-C(=O)NR^{X3A}-$, $-C(=S)NR^{X3A}-$, $-COO-$, $(CHR^{X3A})_k$, $-O-$, $-CH_2NR^{X3A}-$, or $-NR^{X3A}-$, wherein each occurrence of R^{X3A} is independently hydrogen, an aliphatic, heteroaliphatic, aryl, heteroaryl, $-(alkyl)aryl$, $-(alkyl)heteroaryl$, $-(heteroalkyl)aryl$, or $-(heteroalkyl)heteroaryl$ moiety, or R^{X3A} taken together with R^4 may form a cycloaliphatic, cycloheteroaliphatic, aryl or heteroaryl moiety, and k is an integer from 1 to 3.

6. The compound of claim 5, wherein  has one of the following structures:




wherein R^3 , R^4 , R^{X2A} , $R^{X2A'}$, R^{X2B} and t are as defined in claim 5.

7. The compound of claim 5, wherein  has one of the following structures:



wherein R^3 and R^4 are as defined in claim 5.


8. The compound of claim 3 having the structure:



wherein R^{X2A} is hydrogen or an aliphatic, heteroaliphatic, aryl, heteroaryl, -(alkyl)aryl, -(alkyl)heteroaryl, -(heteroalkyl)aryl, or -(heteroalkyl)heteroaryl moiety.

9. The compound of claim 8, wherein R^{X2A} is a substituted or unsubstituted, linear or branched lower alkyl moiety.


10. The compound of claim 8, wherein R^{X2A} is methyl, ethyl, propyl, isopropyl or phenethyl.



wherein each occurrence of R^{2A} is independently hydrogen or lower alkyl; each occurrence of X is independently a halogen; s is an integer from 0 to 3 and u is an integer from 1 to 6; whereby each of the foregoing alkyl moieties may be linear or branched, substituted or unsubstituted and cyclic or acyclic.

20. The compound of claim 19, wherein each occurrence of X is independently chlorine or fluorine and R^{2A} is methyl.

21. The compound of claim 17, wherein R^2 has one of the following structures:



STATEMENT UNDER ARTICLE 19(1)

Applicant respectfully submits that no new matter is presented with the amendments set forth in the "Letter for PCT Article 19 Amendment of Claims" filed concurrently herewith. Specifically, the claims have been amended in order to more clearly set forth what is intended as Applicants' invention, to expedite prosecution or to correct typographical or clerical errors.

For example, claims 3 has been amended (i) to present the claim in independent form, (ii) to add a definition for variable $R^{3'}$, which had been inadvertently omitted at the time of filing; and (iii) to correct some of the functional groups that are recited for variable X^3 (i.e., "-NC(=S)N-, -N-C(=N-C≡N)N-, -NS(O₂)N-" was replaced with "-N(R^{X3A})C(=S)N(R^{X3A})-, -N(R^{X3A})-C(=N-C≡N)N(R^{X3A})-, -N(R^{X3A})S(O₂)N(R^{X3A})-"). Support for the definition of $R^{3'}$ can be found, for example, in original claim 1. With respect to the recitation of functional groups "-NC(=S)N-, -N-C(=N-C≡N)N-, -NS(O₂)N-", amendment is necessary to correct the error in valency on some of the nitrogen atoms. Specifically, the valency in the highlighted nitrogen atoms in -NC(=S)N-, -N-C(=N-C≡N)N- and -NS(O₂)N- is defective. The correct valency is "-N(R^{X3A})C(=S)N(R^{X3A})-, -N(R^{X3A})-C(=N-C≡N)N(R^{X3A})-, -N(R^{X3A})S(O₂)N(R^{X3A})-." Claim 3 has been amended to reflect this. A similar amendment was made in claim 5.

In addition, claim 3, as amended, include provisos (i) – (iii), which specifically exclude compounds disclosed in EP 0 386 611 and EP 0 316 965 (i.e., references cited under category "X" in the International Search Report mailed 20 April 2004). Proviso (ii) finds support in original claim 4 (See proviso (i) in (now canceled) claim 4), and specifically

excludes compounds generically disclosed in EP 0 386 611. Similarly, proviso (i) in amended claim 3 excludes compounds 18-20 disclosed on page 23 of EP 0 316 965; and proviso (iii) excludes compounds generically disclosed in claim 1 of EP 0 316 965.

Claims 6, 7, 10 and 20 were amended to correct claim dependency.

No new matter is introduced by the claim amendments submitted herein. Therefore, Applicant respectfully requests entry of these amendments, and consideration of these amendments in processing the application.





FIG. 1B



INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 03/39276

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07C237/22 C07C237/12 C07C237/14 C07D333/02 C07D307/02 C07D211/04 C07D249/08 C07C311/16 C07C311/08 C07D275/02 C07D285/06 C07D213/04 A61K31/166 A61P25/28 C07K5/02		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07C C07D C07K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, CHEM ABS Data, BEILSTEIN Data, WPI Data, PAJ		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00 56335 A (ELLMAN JONATHAN A ;SKILLMAN A GEOFFREY (US); BI XIAONING (US); HAQ) 28 September 2000 (2000-09-28) claims	1,32-37
X	EP 0 316 965 A (HAESSLE AB) 24 May 1989 (1989-05-24) page 23; examples 18-20 page 32, line 40 -page 33, line 17; claim 1	8-32
A	page 26 -page 27; figures 4,5	33-37
X	EP 0 386 611 A (HOFFMANN LA ROCHE) 12 September 1990 (1990-09-12) claim 1; example 47	8-32
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
6 April 2004		20/04/2004
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel: (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3015		Authorized officer Bedel, C

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 03/39276

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D241/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WOLFE M S ET AL: "SUBSTRATE-BASED DIFLUORO KETONE SELECTIVELY INHIBITS ALZHEIMER'S GAMMA-SECRETASE ACTIVITY" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 41, no. 1, 1998, pages 6-9, XP000938942 ISSN: 0022-2623 the whole document	1,32-37
A	KATERINA BROZKOVA: "Peptidomimetic Inhibitors of Extracellular Aspartic Proteinases of Candida Albicans and Candida Tropicalis" COLLECT.CZECH.CHEM.COMMUN., vol. 64, 1999, pages 130-137, XP009020881 page 133; table I	1,32-37

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

6 April 2004

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bedel, C

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 03/39276

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 35-37
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.: 1-7
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 35-37 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.1

Claims Nos.: 35-37

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

Continuation of Box I.2

Claims Nos.: 1-7

Present claims 1-7 relate to an extremely large number of possible compounds. In fact, the claims contain so many options, variables, possible permutations and provisos that a lack of clarity (and/or conciseness) within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible. Consequently, the search has been carried out for those parts of the application which do appear to be clear (and/or concise), namely claim 8.

Furthermore the initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claim(s) may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of the claims is impossible. Consequently, the search has been restricted to the subject-matter of claim 8 which covers all the examples.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 03/39276

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0056335	A	28-09-2000	AU 3771700 A CA 2367112 A1 EP 1178800 A1 JP 2002539260 T WO 0056335 A1	09-10-2000 28-09-2000 13-02-2002 19-11-2002 28-09-2000
EP 0316965	A	24-05-1989	AU 1516788 A AU 573735 B2 AU 2494484 A DK 478684 A EP 0118223 A1 EP 0316965 A2 ES 8602847 A1 ES 8802129 A1 ES 8802393 A1 ES 8604111 A1 FI 843837 A WO 8403044 A1 NO 843975 A US 4713445 A JP 60500415 T	08-09-1988 23-06-1988 30-08-1984 05-10-1984 12-09-1984 24-05-1989 16-03-1986 16-06-1988 16-08-1988 01-06-1986 28-09-1984 16-08-1984 02-10-1984 15-12-1987 28-03-1985
EP 0386611	A	12-09-1990	AU 634756 B2 AU 5058290 A CA 2010531 A1 CS 9001079 A2 EP 0386611 A2 HU 53067 A2 JP 2279661 A MC 2103 A MX 19616 A NO 901033 A NZ 232695 A PT 93342 A ZA 9001498 A	04-03-1993 06-09-1990 06-09-1990 13-08-1991 12-09-1990 28-09-1990 15-11-1990 11-03-1991 28-02-1994 07-09-1990 25-02-1993 07-11-1990 28-11-1990